PATENT  
Docket No.: 201487/1020 (E2-001PCT-US)

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants	:	Notomi et al.	)	Examiner:
			)	Unknown
Serial No.	:	09/530,061	)	
			)	Art Unit:
Filed	:	April 21, 2000	)	Unknown
			)	
For	:	METHOD OF SYNTHESIZING NUCLEIC	)	
		ACID	)	

Assistant Commissioner for Patents  
Washington, D.C. 20231

Sir:

Transmitted herewith in the above-identified application are:

- ☒ Submission of Combined Declaration and Power of Attorney;
- ☒ Combined Declaration and Power of Attorney; and
- ☒ A self-addressed, prepaid postcard for acknowledging receipt.

- ☒ The Commissioner is hereby authorized to charge any additional fees or credit any overpayment to Deposit Account No. 14-1138.

A duplicate copy of this sheet is enclosed.

Date: August 29, 2000  
Edwin V. Merkel  
Registration No. 40,087Nixon Peabody LLP  
Clinton Square, P.O. Box 31051  
Rochester, New York 14603  
Telephone: (716) 263-1128  
Facsimile: (716) 263-1600

## Certificate of Mailing - 37 CFR 1.8 (a)

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to: Assistant Commissioner for Patents, Washington, D.C. 20231, on the date below.

8/29/00

Date

Wendy L. Harrel

FORM PTO-1390 (Modified)  
(REV 11-98)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

## TRANSMITTAL LETTER TO THE UNITED STATES

201487/1020

DESIGNATED/ELECTED OFFICE (DO/EO/US)

U.S. APPLICATION NO. (IF KNOWN) 37 CFR

CONCERNING A FILING UNDER 35 U.S.C. 371

01 Rec'd PCT/PTO 13 OCT 2000

INTERNATIONAL APPLICATION NO.  
PCT/JP99/06213INTERNATIONAL FILING DATE  
08 November 1999PRIORITY DATE CLAIMED  
09 November 1999

09/53 U061

TITLE OF INVENTION

METHOD OF SYNTHESIZING NUCLEIC ACID

APPLICANT(S) FOR DO/EO/US

NOTOMI, Tsugunori and HASE, Tetsu

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☐ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
  - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☒ has been transmitted by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ A copy of the International Search Report (PCT/ISA/210).
8. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
  - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
9. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
10. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
11. ☐ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).

Items 13 to 20 below concern document(s) or information included:

13. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☒ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☒ Certificate of Mailing by Express Mail
20. ☒ Other items or information:

- 1) **UNSIGNED** Combined Declaration and Power of Attorney.
- 2) Statement re sequence listing with attached diskette.

\*\*\*\*\*  
\*\*\* RX REPORT \*\*\*  
\*\*\*\*\*

RECEPTION OK

TX/RX NO	6558	
CONNECTION TEL		716 263 1600
SUBADDRESS		
CONNECTION ID		
ST. TIME	10/13 16:27	
USAGE T	00'34	
PGS.	2	
RESULT	OK	

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant(s) :	Tsugunori Notomi and Tetsu Hase	)	Examiner:
		)	To Be Assigned
Serial No. :	To Be Assigned (based on PCT/JP99/06213 filed	)	
	November 8, 1999)	)	Art Unit:
		)	To Be Assigned
Filed :	Herewith	)	
		)	
For :	METHOD OF SYNTHESIZING NUCLEIC	)	
	ACID	)	

PRELIMINARY AMENDMENT

Commissioner of Patents and Trademarks  
Washington, D.C. 20231  
**Box: PCT**

Dear Sir:

Please amend the above-identified patent application as follows:

In the Specification:

On page 26, line 22, please change "P" to -- O --.

In the Claims:

Please delete claim 12 and add new claims 25-28 as follows:

--25. The method according to claim 5, wherein the step of rendering base pairing ready in steps iii) and iv) is conducted by the strand displacement synthesis of complementary chain by a polymerase catalyzing the strand displacement reaction of synthesizing complementary chain wherein an outer primer annealing to the 3'-side of F2c in the template and an outer primer annealing to the 3'-side of the region used as the origin of synthesis in step iv) for the first nucleic acid serve as the origin of synthesis.--

--26. The method according to claim 9, wherein the strand displacement reaction of synthesizing complementary chain is carried out in the presence of a melting temperature regulator.--

--27. The method according to claim 26, wherein the melting temperature regulator is betaine.--

--28. The method according to claim 27, wherein 0.2 to 3.0 M betaine is allowed to be present in the reaction solution.--

Please amend claims 3-8, 13, 16, 19, 22, and 24 as follows:

3. (Amended) An oligonucleotide [composed of] comprising at least two regions X2 and X1c below, and X1c is linked to the 5'-side of X2,

X2: a region having a nucleotide sequence complementary to an arbitrary region X2c in nucleic acid having a specific nucleotide sequence, and

X1c: a region having substantially the same nucleotide sequence as in a region X1c located at the 5'-side of the region X2c in nucleic acid having a specific nucleotide sequence.

4. (Amended) The method according to claim 1, wherein the nucleic acid in step a) is second nucleic acid provided by the following steps:

i) the step of annealing, to a region F2c in nucleic acid serving as a template, of a region F2 in [the] an oligonucleotide [described in claim 3 wherein the region X2 is a region F2 and the region X1c is a region F1c] comprising at least two regions below, the region F2 and a region F1c linked to the 5'-side of F2,

F2: a region having a nucleotide sequence complementary to an arbitrary region F2c in nucleic acid having a specific nucleotide sequence, and

F1c: a region having substantially the same nucleotide sequence as in a region F1c located at the 5'-side of the region F2c in nucleic acid having a specific nucleotide sequence,

ii) the step of synthesizing first nucleic acid having a nucleotide sequence complementary to the template wherein F2 in the oligonucleotide serves as the origin of synthesis,

iii) the step of rendering an arbitrary region in the first nucleic acid synthesized in step ii) ready for base pairing, and

iv) the step of annealing an oligonucleotide having a nucleotide sequence complementary to the region made ready for base pairing in the first nucleic acid in step iii), followed by synthesizing second nucleic acid with said oligonucleotide as the origin of synthesis and rendering F1 at the 3'-terminal thereof ready for base pairing.

5. (Amended) The method according to claim 4, wherein the region enabling base pairing in step iii) is R2c, and the oligonucleotide in step iv) is [the] an oligonucleotide [described in claim 3 wherein the region X2c is a region R2c and the region X1c is a region R1c] comprising at least two regions below, the region R2 and a region R1c linked to the 5'-side of R2,

R2: a region having a nucleotide sequence complementary to an arbitrary region R2c in nucleic acid having a specific nucleotide sequence, and

R1c: a region having substantially the same nucleotide sequence as in a region R1c located at the 5'-side of the region R2c in nucleic acid having a specific nucleotide sequence.

6. (Amended) The method according to claim 4 [or 5], wherein the step of rendering base pairing ready in steps iii) and iv) is conducted by the strand displacement synthesis of complementary chain by a polymerase catalyzing the strand displacement reaction of synthesizing complementary chain wherein an outer primer annealing to the 3'-side of F2c in the template and an outer primer annealing to the 3'-side of the region used as the origin of synthesis in step iv) for the first nucleic acid serve as the origin of synthesis.

7. (Amended) The method according to claim [6] 25, wherein the melting temperature of each oligonucleotide and its complementary region in the template used in the reaction is in the following relationship under the same stringency:

(outer primer/region at the 3'-side in the template)  $\leq$  (F2c/F2 and R2c/R2)  $\leq$  (F1c/F1 and R1c/R1).

8. (Amended) The method according to [any one of claims 4 to 7] claim 4, wherein the nucleic acid serving as the template is RNA, and the synthesis of complementary chain in step ii) is conducted by an enzyme having a reverse transcriptase activity.

13. (Amended) The method according to claim 1 [or 9], wherein the strand displacement reaction of synthesizing complementary chain is carried out in the presence of a melting temperature regulator.

16. (Amended) A method of detecting a target nucleotide sequence in a sample, which comprises performing an amplification method [described in any one of claims 9 to 15] according to claim 9 and observing whether an amplification reaction product is generated or not.

19. (Amended) The method according to claim 16, wherein [an] the amplification method [described in any one of claims 9 to 15] is conducted in the presence of a detector for nucleic acid, and whether an amplification reaction product is generated or not is observed on the basis of a change in the signal of the detector.

22. (Amended) The kit according to claim 21, wherein the oligonucleotide in ii) is [the] an oligonucleotide [described in claim 3] comprising at least two regions X2 and X1c below, and X1c is linked to the 5'-side of X2.

X2: a region having a nucleotide sequence complementary to an arbitrary region X2c in nucleic acid having a specific nucleotide sequence, and


X1c: a region having substantially the same nucleotide sequence as in a region X1c located at the 5'-side of the region X2c in nucleic acid having a specific nucleotide sequence

wherein an arbitrary region R2c in a complementary chain synthesized with the oligonucleotide in i) as the origin of synthesis is X2c, and R1c located at the 5' of R2c is X1c.

24. (Amended) A kit for detection of a target nucleotide sequence, comprising a detector for detection of a product of nucleic acid synthetic reaction additionally in a kit described in [any one of claims 21 to 23] claim 21.

In view of the foregoing preliminary amendments, examination of the present application is respectfully requested.

Respectfully submitted,



---

Edwin V. Merkel  
Registration No. 40,087

Date: April 21, 2000

NIXON PEABODY LLP  
Clinton Square, P.O. Box 1051  
Rochester, New York 14603  
Telephone: (716) 263-1128  
Facsimile: (716) 263-1600



EXPRESS MAIL CERTIFICATE

DOCKET NO.: 201487/1020

APPLICANT(S): Notomi et al.

TITLE: METHOD OF SYNTHESIZING NUCLEIC ACID


Certificate is attached to the Statement re sequence listing  
w/diskette) of the above-named application.

"EXPRESS MAIL" NUMBER: EL434485635US

DATE OF DEPOSIT: April 21, 2000

I hereby certify that this paper or fee is being  
deposited with the United States Postal Service "Express Mail Post  
Office to Addressee" service under 37 CFR 1.10 on the date  
indicated above and is addressed to the Commissioner of Patents and  
Trademarks, Box PCT, Washington, D.C. 20231.

Sherri A. Moscato  
(Typed or printed name of person  
mailing paper or fee)

  
(Signature of person mailing paper or  
fee)

PATENT  
Docket No.: 201487/1020 (E2-001PCT-US)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants	:	Tsugunori Notomi and Tetsu Hase	)	Examiner:
			)	To Be Assigned
Serial No.	:	To Be Assigned (based on PCT/JP99/06213 filed	)	
		November 8, 1999)	)	Art Unit:
			)	To Be Assigned
Filed	:	Herewith	)	
			)	
For	:	METHOD OF SYNTHESIZING NUCLEIC	)	
		ACID	)	

---

STATEMENT IN ACCORDANCE WITH 37 CFR § 1.821(f)


Commissioner of Patent and Trademarks  
Washington, D.C. 20231  
**Box: PCT**

Sir:

In accordance with 37 CFR § 1.821(f), this statement confirms that the contents of the Sequence Listing attached to the subject application as submitted herewith and on the computer readable 3.5" Diskette submitted herewith are the same.

Respectfully submitted,

Date: Apr. 1 21, 2000

  
Edwin V. Merkel  
Registration No. 40,087

NIXON PEABODY LLP  
Clinton Square, P.O. Box 1051  
Rochester, New York 14603  
Telephone: (716) 263-1128  
Facsimile: (716) 263-1600

18/PRTS

526 Rec'd PCT/ITC 21 APR 2000

## DESCRIPTION

## METHOD OF SYNTHESIZING NUCLEIC ACID

## Technical Field

The present invention relates to a method of synthesizing nucleic acid composed of a specific nucleotide sequence, which is useful as a method of amplifying nucleic acid.

## Background Art

An analysis method based on complementarity of a nucleic acid nucleotide sequence can analyze genetic traits directly. Accordingly, this analysis is a very powerful means for identification of genetic diseases, canceration, microorganisms etc. Further, a gene itself is the object of detection, and thus time-consuming and cumbersome procedures such as in culture can be omitted in some cases.

Nevertheless, the detection of a target gene present in a very small amount in a sample is not easy in general so that amplification of a target gene itself or its detection signal is necessary. As a method of amplifying a target gene, the PCR (polymerase chain reaction) method is known (Science, 230, 1350-1354, 1985). Currently, the PCR method is the most popular method as a technique of amplifying nucleic acid in vitro. This method was established firmly as an excellent detection method by virtue of high sensitivity based on the effect of exponential

amplification. Further, since the amplification product can be recovered as DNA, this method is applied widely as an important tool supporting genetic engineering techniques such as gene cloning and structural determination. In the PCR method, however, there are the following noted problems: a special temperature controller is necessary for practice; the exponential progress of the amplification reaction causes a problem in quantification; and samples and reaction solutions are easily contaminated from the outside to permit nucleic acid mixed in error to function as a template.

As genomic information is accumulated, analysis of single nucleotide polymorphism (SNPs) comes to attract attention. Detection of SNPs by means of PCR is feasible by designing a primer such that its nucleotide sequence contains SNPs. That is, whether a nucleotide sequence complementary to the primer is present or not can be inferred by determining whether a reaction product is present or not. However, once a complementary chain is synthesized in error in PCR by any chance, this product functions as a template in subsequent reaction, thus causing an erroneous result. In practice, it is said that strict control of PCR is difficult with the difference of only one base given at the terminal of the primer. Accordingly, it is necessary to improve specificity in order to apply PCR to detection of SNPs.

On one hand, a method of synthesizing nucleic acid by a

ligase is also practically used. The LCR method (ligase chain reaction, Laffler TG; Garrino JJ; Marshall RL; Ann. Biol. Clin. (Paris), 51:9, 821-6, 1993) is based on the reaction in which two adjacent probes are hybridized with a target sequence and ligated to each other by a ligase. The two probes could not be ligated in the absence of the target nucleotide sequence, and thus the presence of the ligated product is indicative of the target nucleotide sequence. Because the LCR method also requires control of temperature for separation of a complementary chain from a template, there arises the same problem as in the PCR method. For LCR, there is also a report on a method of improving specificity by adding the step of providing a gap between adjacent probes and filling the gap by a DNA polymerase. However, what can be expected in this modified method is specificity only, and there still remains a problem in that control of temperature is required. Furthermore, use of the additional enzyme leads to an increase in cost.

A method called the SDA method (strand displacement amplification) [Proc. Natl. Acad. Sci. USA, 89, 392-396, 1992] [Nucleic Acid. Res., 20, 1691-1696, 1992] is also known as a method of amplifying DNA having a sequence complementary to a target sequence as a template. In the SDA method, a special DNA polymerase is used to synthesize a complementary chain starting from a primer complementary to the 3'-side of a certain

nucleotide sequence while displacing a double-stranded chain if any at the 5'-side of the sequence. In the present specification, the simple expression "5'-side" or "3'-side" refers to that of a chain serving as a template. Because a double-stranded chain at the 5'-side is displaced by a newly synthesized complementary chain, this technique is called the SDA method. The temperature-changing step essential in the PCR method can be eliminated in the SDA method by previously inserting a restriction enzyme recognition sequence into an annealed sequence as a primer. That is, a nick generated by a restriction enzyme gives a 3'-OH group acting as the origin of synthesis of complementary chain, and the previously synthesized complementary chain is released as a single-stranded chain by strand displacement synthesis and then utilized again as a template for subsequent synthesis of complementary chain. In this manner, the complicated control of temperature essential in the PCR method is not required in the SDA method.

In the SDA method, however, the restriction enzyme generating a nick should be used in addition to the strand displacement-type DNA polymerase. This requirement for the additional enzyme is a major cause for higher cost. Further, because the restriction enzyme is to be used not for cleavage of both double-stranded chains but for introduction of a nick (that is, cleavage of only one of the chains), a dNTP derivative

such as  $\alpha$ -thio dNTP should be used as a substrate for synthesis to render the other chain resistant to digestion with the enzyme. Accordingly, the amplification product by SDA has a different structure from that of natural nucleic acid, and there is a limit to cleavage with restriction enzymes or application of the amplification product to gene cloning. In this respect too, there is a major cause for higher cost. In addition, when the SDA method is applied to an unknown sequence, there is the possibility that the same nucleotide sequence as the restriction enzyme recognition sequence used for introducing a nick may be present in a region to be synthesized. In this case, it is possible that a complete complementary chain is prevented from being synthesized.

NASBA (nucleic acid sequence-based amplification, also called the TMA/transcription mediated amplification method) is known as a method of amplifying nucleic acid wherein the complicated control of temperature is not necessary. NASBA is a reaction system wherein DNA is synthesized by DNA polymerase in the presence of target RNA as a template with a probe having a T7 promoter added thereto, and the product is formed with a second probe into a double-stranded chain, followed by transcription by T7 RNA polymerase with the formed double-stranded chain as a template to amplify a large amount of RNA (Nature, 350, 91-92, 1991). NASBA requires some heat denaturation steps until double-stranded DNA is completed, but

the subsequent transcriptional reaction by T7 RNA polymerase proceeds under isothermal conditions. However, a combination of plural enzymes such as reverse transcriptase, RNase H, DNA polymerase and T7 RNA polymerase is essential, and this is unfavorable for cost similarly to SDA. Further, because it is complicated to set up conditions for a plurality of enzyme reaction, this method is hardly widespread as a general analytical method. In the known reactions of amplification of nucleic acid, there remain problems such as complicated control of temperature and the necessity for plural enzymes as described above

For these known reactions of synthesizing nucleic acid, there are few reports on an attempt for further improving the efficiency of synthesis of nucleic acid without sacrificing specificity or cost. For example, in a method called RCA (rolling-circle amplification), it was shown that single-stranded DNA having a series of nucleotide sequences complementary to a padlock probe can be synthesized continuously in the presence of a target nucleotide sequence (Paul M. Lizardi et al., *Nature Genetics*, 19, 225-232, July, 1998). In RCA, a padlock probe having a special structure wherein each of the 5'- and 3'-terminals of a single oligonucleotide constitutes an adjacent probe in LCR is utilized. Then, the continuous reaction of synthesizing complementary chain with the padlock probe as a template which



was ligated and cyclized in the presence of a target nucleotide sequence is triggered by combination with a polymerase catalyzing the strand displacement-type reaction of synthesizing complementary chain. Single-stranded nucleic acid having a structure of a series of regions each consisting of the same nucleotide sequence is thus formed. A primer is further annealed to this single-stranded nucleic acid to synthesize its complementary chain and a high degree of amplification is thus realized. However, there still remains the problem of the necessity for a plurality of enzymes. Further, triggering of synthesis of complementary chain depends on the reaction of ligating two adjacent regions, and its specificity is basically the same as in LCR.

For the object of supplying 3'-OH, there is a known method in which a nucleotide sequence is provided at the 3'-terminal with a sequence complementary thereto and a hair pin loop is formed at the terminal (Gene, 71, 29-40, 1988). Synthesis of complementary chain with a target sequence itself as a template starts at the hairpin loop to form single-stranded nucleic acid composed of the complementary nucleotide sequence. For example, a structure in which annealing occurs in the same chain at the terminal to which a complementary nucleotide sequence has been linked is realized in PCT/FR95/00891. In this method, however, the step in which the terminal cancels base pairing with the complementary chain and base pairing is constituted again in

the same chain is essential. It is estimated that this step proceeds depending on a subtle equilibrium state at the terminal of mutually complementary nucleotide sequences involving base pairing. That is, an equilibrium state maintained between base pairing with a complementary chain and base pairing in the same chain is utilized and the only chain annealing to the nucleotide sequence in the same chain serves as the origin of synthesis of a complementary chain. Accordingly, it is considered that strict reaction conditions should be set to achieve high reaction efficiency. Further, in this prior art, the primer itself forms a loop structure. Accordingly, once a primer dimer is formed, amplification reaction is automatically initiated regardless of whether a target nucleotide sequence is present or not, and an unspecific synthetic product is thus formed. This can be a serious problem. Further, formation of the primer dimer and subsequent consumption of the primer by unspecific synthetic reaction lead to a reduction in the amplification efficiency of the desired reaction.

Besides, there is a report that a region not serving as a template for DNA polymerase was utilized to realize a 3'-terminal structure annealing to the same chain (EP713922). This report also has the same problem as in PCT/FR95/00891 supra in respect of the utilization of dynamic equilibrium at the terminal or the possibility of unspecific synthetic reaction due to formation of a dimer primer. Further, a special region

not serving as a template for DNA polymerase should be prepared as a primer.

Further, in various signal amplification reactions to which the principle of NASBA described above is applied, an oligonucleotide having a hairpin structure at the terminal thereof is often utilized to supply a double-stranded promoter region (JP-A 5-211873). However, these techniques are not those permitting successive supply of 3'-OH for synthesis of a complementary chain. Further, a hairpin loop structure having a 3'-terminal annealed in the same chain is utilized for the purpose of obtaining a DNA template transcribed by RNA polymerase is utilized in JP-A 10-510161 (WO96/17079). In this method, the template is amplified by using transcription into RNA and reverse transcription from RNA to DNA. In this method, however, the reaction system cannot be constituted without a combination of a plurality of enzymes.

#### Disclosure of the Invention

The object of the present invention is to provide a method of synthesizing nucleic acid based on a novel principle. A more specific object is to provide a method capable of realizing the synthesis of nucleic acid depending on sequence efficiently at low costs. That is, an object of the present invention is to provide a method capable of achieving the synthesis and amplification of nucleic acid by a single enzyme even under

isothermal reaction conditions. Another object of the present invention is to provide a method of synthesizing nucleic acid which can realize high specificity difficult to achieve in the known reaction principle of nucleic acid synthesis, as well as a method of amplifying nucleic acid by applying said synthetic method.

The present inventors focused their attention on the fact that the utilization of a polymerase catalyzing strand displacement-type synthesis of complementary chain is useful for nucleic acid synthesis not depending on complicated control of temperature. Such a DNA polymerase is an enzyme utilized in SDA and RCA. However, even if such an enzyme is used, another enzyme reaction is always required for supplying 3'-OH as the origin of synthesis in the known means based on primers, such as SDA.

Under these circumstances, the present inventors examined supply of 3'-OH from a completely different viewpoint from the known approach. As a result, the present inventors found that by utilizing an oligonucleotide having a special structure, 3'-OH can be supplied without any additional enzyme reaction, thereby completing the present invention. That is, the present invention relates to a method of synthesizing nucleic acid, a method of amplifying nucleic acid by applying said method of synthesizing nucleic acid and a novel oligonucleotide enabling said methods, as follows:

1. A method of synthesizing nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain, comprising:

a) the step of giving nucleic acid which is provided at the 3'-terminal thereof with a region F1 capable of annealing to a part F1c in the same chain and which upon annealing of the region F1 to F1c, is capable of forming a loop containing a region F2c capable of base pairing,

b) the step of performing synthesis of a complementary chain wherein the 3'-terminal of F1 having annealed to F1c serves as the origin of synthesis,

c) the step of annealing, to a region F2c, of an oligonucleotide provided with the 3'-terminal thereof with F2 consisting of a sequence complementary to the region F2c, followed by synthesis, with said oligonucleotide as the origin of synthesis, of a complementary chain by a polymerase catalyzing the strand displacement reaction of synthesizing a complementary chain to displace the complementary chain synthesized in step b), and

d) the step of annealing, to the complementary chain displaced in step c) to be ready for base pairing, of a polynucleotide provided at the 3'-terminal thereof with a sequence complementary to an arbitrary region in said chain synthesized

in step c), followed by synthesis, with said 3'-terminal as the origin of synthesis, of a complementary chain by a polymerase catalyzing the strand displacement reaction of synthesizing a complementary chain to displace the complementary chain synthesized in step c).

2. The method according to item 1, wherein in step d), the origin of synthesis is a region R1 present at the 3'-terminal in the same chain and capable of annealing to a region R1c, and a loop containing the region R2c capable of base pairing is formed by annealing R1 to R1c.

3. An oligonucleotide composed of at least two regions X2 and X1c below, and X1c is linked to the 5'-side of X2,

X2: a region having a nucleotide sequence complementary to an arbitrary region X2c in nucleic acid having a specific nucleotide sequence, and

X1c: a region having substantially the same nucleotide sequence as in a region X1c located at the 5'-side of the region X2c in nucleic acid having a specific nucleotide sequence.

4. The method according to item 1, wherein the nucleic acid in step a) is second nucleic acid provided by the following steps:

i) the step of annealing, to a region F2c in nucleic acid serving as a template, of a region F2 in the oligonucleotide described in item 3 wherein the region X2 is a region F2 and the region X1c is a region F1c,

ii) the step of synthesizing first nucleic acid having a

nucleotide sequence complementary to the template wherein F2 in the oligonucleotide serves as the origin of synthesis, iii) the step of rendering an arbitrary region in the first nucleic acid synthesized in step ii) ready for base pairing, and

iv) the step of annealing an oligonucleotide having a nucleotide sequence complementary to the region made ready for base pairing in the first nucleic acid in step iii), followed by synthesizing second nucleic acid with said oligonucleotide as the origin of synthesis and rendering F1 at the 3'-terminal thereof ready for base pairing.

5. The method according to item 4, wherein the region enabling base pairing in step iii) is R2c, and the oligonucleotide in step iv) is the oligonucleotide described in item 3 wherein the region X2c is a region R2c and the region X1c is a region R1c.

6. The method according to item 4 or 5, wherein the step of rendering base pairing ready in steps iii) and iv) is conducted by the strand displacement synthesis of complementary chain by a polymerase catalyzing the strand displacement reaction of synthesizing complementary chain wherein an outer primer annealing to the 3'-side of F2c in the template and an outer primer annealing to the 3'-side of the region used as the origin of synthesis in step iv) for the first nucleic acid serve as the origin of synthesis.

7. The method according to item 6, wherein the melting

temperature of each oligonucleotide and its complementary region in the template used in the reaction is in the following relationship under the same stringency:

(outer primer/region at the 3'-side in the template)  $\leq$  (F2c/F2 and R2c/R2)  $\leq$  (F1c/F1 and R1c/R1).

8. The method according to any one of items 4 to 7, wherein the nucleic acid serving as the template is RNA, and the synthesis of complementary chain in step ii) is conducted by an enzyme having a reverse transcriptase activity.

9. A method of amplifying nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain by repeatedly conducting the following steps:

A) the step of providing a template which is provided at the 3'- and 5'-terminals thereof with a region consisting of a nucleotide sequence complementary to each terminal region in the same chain and which upon annealing of these mutually complementary nucleotide sequences, forms a loop capable of base pairing therebetween,

B) the step of performing the synthesis of complementary chain wherein the 3'-terminal of said template annealed to the same chain serves as the origin of synthesis,

C) the step of annealing, to the loop portion, of an oligonucleotide provided at the 3'-terminal thereof with a complementary nucleotide sequence to a loop which among said loops, is located at the 3'-terminal site, followed by synthesis,



with the oligonucleotide as the origin of synthesis, of a complementary chain by a polymerase catalyzing the strand displacement reaction of synthesizing a complementary chain to displace the complementary chain synthesized in step B) to make the 3'-terminal thereof ready for base pairing, and

D) the step wherein the chain with the 3'-terminal made ready for base pairing in step C) serves as a new template.

10. The method according to item 9, wherein the oligonucleotide in step C) is provided at the 5'-terminal thereof with a nucleotide sequence complementary to the 3'-terminal serving as the origin of synthesis in step B).

11. The method according to item 10, further comprising the step where a complementary chain synthesized with the oligonucleotide in step C) as the origin of synthesis is used as a template in step A).

12. The method according to item 9, wherein the template in step A) is synthesized by the method described in item 5.

13. The method according to item 1 or 9, wherein the strand displacement reaction of synthesizing complementary chain is carried out in the presence of a melting temperature regulator.

14. The method according to item 13, wherein the melting temperature regulator is betaine.

15. The method according to item 14, wherein 0.2 to 3.0 M betaine is allowed to be present in the reaction solution.

16. A method of detecting a target nucleotide sequence in a

sample, which comprises performing an amplification method described in any one of items 9 to 15 and observing whether an amplification reaction product is generated or not.

17. The method according to item 16, wherein a probe containing a nucleotide sequence complementary to the loop is added to the amplification reaction product and hybridization therebetween is observed.

18. The method according to item 17, wherein the probe is labeled on particles and aggregation reaction occurring upon hybridization is observed.

19. The method according to item 16, wherein an amplification method described in any one of items 9 to 15 is conducted in the presence of a detector for nucleic acid, and whether an amplification reaction product is generated or not is observed on the basis of a change in the signal of the detector.

20. A method of detecting a mutation in a target nucleotide sequence by the detection method described in item 16, wherein a mutation in a nucleotide sequence as the subject of amplification prevents synthesis of any one of complementary chains constituting the amplification method.

21. A kit for synthesis of nucleic acid having complementary chains alternately linked in a single-stranded chain, comprising the following elements:

i) the oligonucleotide described in item 3 wherein the region F2c in nucleic acid as a template is X2c, and F1c located at

the 5'-side of F2c is X1c;

ii) an oligonucleotide containing a nucleotide sequence complementary to an arbitrary region in a complementary chain synthesized with the oligonucleotide in (i) as a primer;

iii) an oligonucleotide having a nucleotide sequence complementary to a region F3c located at the 3'-side of the region F2c in the nucleic acid serving as a template;

iv) a DNA polymerase catalyzing the strand displacement-type reaction of synthesizing complementary chain; and

v) a nucleotide serving as a substrate for the element iv).

22. The kit according to item 21, wherein the oligonucleotide in ii) is the oligonucleotide described in item 3 wherein an arbitrary region R2c in a complementary chain synthesized with the oligonucleotide in i) as the origin of synthesis is X2c, and R1c located at the 5' of R2c is X1c.

23. The kit according to item 22, further comprising:

vi) an oligonucleotide having a nucleotide sequence complementary to a region R3c located at the 3'-side of the arbitrary R2c in a complementary chain synthesized with the oligonucleotide in i) as the origin of synthesis.

24. A kit for detection of a target nucleotide sequence, comprising a detector for detection of a product of nucleic acid synthetic reaction additionally in a kit described in any one of items 21 to 23.

The nucleic acid having complementary nucleotide

sequences linked alternately in a single-stranded chain as the object of synthesis in the present invention means nucleic acid having mutually complementary nucleotide sequences linked side by side in a single-stranded chain. Further, in the present invention, it should contain a nucleotide sequence for forming a loop between the complementary chains. In the present invention, this sequence is called the loop-forming sequence. The nucleic acid synthesized by the present invention is composed substantially of mutually complementary chains linked via the loop-forming sequence. In general, a strand not separated into 2 or more molecules upon dissociation of base pairing is called a single-stranded chain regardless of whether it partially involves base pairing or not. The complementary nucleotide sequence can form base pairing in the same chain. An intramolecular base-paired product, which can be obtained by permitting the nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain according to the present invention to be base-paired in the same chain, gives a region constituting an apparently double-stranded chain and a loop not involving base pairing.

That is, the nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain according to the present invention contains complementary nucleotide sequences capable of annealing in the same chain, and its annealed product can be defined as single-stranded

nucleic acid constituting a loop not involving base pairing at a bent hinged portion. A nucleotide having a nucleotide sequence complementary thereto can anneal to the loop not involving base pairing. The loop-forming sequence can be an arbitrary nucleotide sequence. The loop-forming sequence is capable of base pairing so as to initiate the synthesis of a complementary chain for displacement, and is provided preferably with a sequence distinguishable from a nucleotide sequence located in the other region in order to achieve specific annealing. For example, in a preferred embodiment, the loop-forming sequence contains substantially the same nucleotide sequence as a region F2c (or R2c) located at the 3'-side of a region (i.e. F1c or R1c) derived from nucleic acid as a template and annealed in the same chain.

In the present invention, substantially the same nucleotide sequence is defined as follows. That is, when a complementary chain synthesized with a certain sequence as a template anneals to a target nucleotide sequence to give the origin of synthesizing a complementary chain, this certain sequence is substantially the same as the target nucleotide sequence. For example, substantially the same sequence as F2 includes not only absolutely the same nucleotide sequence as F2 but also a nucleotide sequence capable of functioning as a template giving a nucleotide sequence capable of annealing to F2 and acting as the origin of synthesizing complementary chain.

The term "anneal" in the present invention means formation of a double-stranded structure of nucleic acid through base pairing based on the law of Watson-Crick. Accordingly, even if a nucleic acid chain constituting base pairing is a single-stranded chain, annealing occurs if intramolecular complementary nucleotide sequences are base-paired. In the present invention, annealing and hybridization have the same meaning in that the nucleic acid constitutes a double-stranded structure through base pairing.

The number of pairs of complementary nucleotide sequences constituting the nucleic acid according to the present invention is at least 1. According to a desired mode of the present invention, it may be 2 or more. In this case, there is theoretically no upper limit to the number of pairs of complementary nucleotide sequences constituting the nucleic acid. When the nucleic acid as the synthetic product of the present invention is constituted of plural sets of complementary nucleotide sequences, this nucleic acid is composed of repeated identical nucleotide sequences.

The nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain synthesized by the present invention may not have the same structure as naturally occurring nucleic acid. It is known that if a nucleotide derivative is used as a substrate when nucleic acid is synthesized by the action of a DNA polymerase, a nucleic

acid derivative can be synthesized. The nucleotide derivative used includes nucleotides labeled with a radioisotope or nucleotide derivatives labeled with a binding ligand such as biotin or digoxin. These nucleotide derivatives can be used to label nucleic acid derivatives as the product. Alternatively, if fluorescent nucleotides are used as a substrate, the nucleic acid as the product can be a fluorescent derivative. Further, this product may be either DNA or RNA. Which one is formed is determined by a combination of the structure of a primer, the type of a substrate for polymerization and polymerization reagents for carrying out polymerization of nucleic acid.

Synthesis of the nucleic acid having the structure described above can be initiated by use of a DNA polymerase having the strand displacement activity and nucleic acid which is provided at the 3'-terminal thereof with a region F1 capable of annealing to a part F1c in the same chain and which upon annealing of the region F1 to F1c, is capable of forming a loop containing a region F2c capable of base pairing. There are many reports on the reaction of synthesizing complementary chain wherein a hairpin loop is formed and a sample sequence itself is used as a template, while in the present invention the portion of the hairpin loop is provided with a region capable of base pairing, and there is a novel feature on utilization of this region in synthesizing complementary chain. By use of this

region as the origin of synthesis, a complementary chain previously synthesized with a sample sequence itself as a template is displaced. Then, a region R1c (arbitrary region) located at the 3'-terminal of the displaced chain is in a state ready for base-pairing. A region having a complementary sequence to this R1c is annealed thereto, resulting in formation of the nucleic acid (2 molecules) having a nucleotide sequence extending from F1 to R1c and its complementary chain linked alternately via the loop-forming sequence. In the present invention, the arbitrary region such as R1c above can be selected arbitrarily provided that it can be annealed to a polynucleotide having a nucleotide sequence complementary to that region, and that a complementary chain synthesized with the polynucleotide as the origin of synthesis has necessary functions for the present invention.

In the present invention, the term "nucleic acid" is used. The nucleic acid in the present invention generally includes both DNA and RNA. However, nucleic acid whose nucleotide is replaced by an artificial derivative or modified nucleic acid from natural DNA or RNA is also included in the nucleic acid of the present invention insofar as it functions as a template for synthesis of complementary chain. The nucleic acid of the present invention is generally contained in a biological sample. The biological sample includes animal, plant or microbial tissues, cells, cultures and excretions, or extracts therefrom.



The biological sample of the present invention includes intracellular parasitic genomic DNA or RNA such as virus or mycoplasma. The nucleic acid of the present invention may be derived from nucleic acid contained in said biological sample. For example, cDNA synthesized from mRNA, or nucleic acid amplified on the basis of nucleic acid derived from the biological sample, is a typical example of the nucleic acid of the present invention.

The nucleic acid characteristic of the present invention which is provided at the 3'-terminal thereof with a region F1 capable of annealing to a part F1c in the same chain and which upon annealing of the region F1 to F1c, is capable of forming a loop containing a region F2c capable of base pairing can be obtained in various methods. In the most preferable embodiment, the reaction of synthesizing complementary chain utilizing an oligonucleotide having the following structure can be used to give the structure.

That is, the useful oligonucleotide in the present invention consists of at least two regions X2 and X1c below wherein X1c is ligated to the 5'-side of X2.

X2: a region having a nucleotide sequence complementary to a region X2c in nucleic acid having a specific nucleotide sequence.

X1c: a region having substantially the same nucleotide sequence as a region X1c located at the 5'-side of the region

X2c in nucleic acid having a specific nucleotide sequence.

Here, the nucleic acid having a specific nucleotide sequence by which the structure of the oligonucleotide of the invention is determined refers to nucleic acid serving as a template when the oligonucleotide of the present invention is used as a primer. In the case of detection of nucleic acid based on the synthetic method of the present invention, the nucleic acid having a specific nucleotide sequence is a detection target or nucleic acid derived from the detection target. The nucleic acid having a specific nucleotide sequence refers to nucleic acid wherein at least a part of the nucleotide sequence is revealed or predictable. The part of the nucleotide sequence revealed is the region X2c and the region X1c located at the 5'-side thereof. It can be supposed that these 2 regions are contiguous or located apart from each other. By the relative positional relationship of the two, the state of a loop formed upon self-annealing of nucleic acid as the product is determined. The distance between the two is preferably not very apart from each other in order that nucleic acid as the product is subjected to self-annealing preferentially over intermolecular annealing. Accordingly, the positional relationship of the two is preferably that they are contiguous via a distance of usually 0 to 100 bases. However, in the formation of a loop by self-annealing described below, there can be the case where it would be disadvantageous for formation of a loop in a desired

state that the two are too close to each other. In the loop, there is a need for a structure for annealing of a new oligonucleotide and for readily initiating the strand-displacement reaction of synthesizing a complementary chain with said oligonucleotide as the origin of synthesis. More preferably, the distance between the region X2c and the region X1c located at the 5'-side of X2c is designed to be 0 to 100 bases, more desirably 10 to 70 bases. This numerical value shows a length excluding X1c and X2. The number of bases constituting the part of a loop is that of this length plus a region corresponding to X2.

Both the terms "same" and "complementary" used for characterization of the nucleotide sequence constituting the oligonucleotide based on the present invention do not mean being absolutely the same or absolutely complementary. That is, the same sequence as a certain sequence includes sequences complementary to nucleotide sequences capable of annealing to a certain sequence. On the other hand, the complementary sequence means a sequence capable of annealing under stringent conditions to provide a 3'-terminal serving as the origin of synthesis of complementary chain.

Usually, the regions X2 and X1c constituting the oligonucleotide of the present invention for the nucleic acid having a specific nucleotide sequence are located contiguously without being overlapped. If there is a common part in both

the nucleotide sequences, the two can be partially overlaid. Because X2 should function as a primer, it should always be a 3'-terminal. On the other hand, X1c should give the function of a primer as described below to the 3'-terminal of a complementary chain synthesized with the nucleic acid as a template, and thus it shall be arranged at the 5'-terminal. The complementary chain obtained with this oligonucleotide as the origin of synthesis serves as a template for synthesis of complementary chain in the reverse direction in the next step, and finally the part of the oligonucleotide of the present invention is copied as a template into a complementary chain. The 3'-terminal generated by copying has the nucleotide sequence X1, which anneals to X1c in the same chain to form a loop.

In the present invention, the oligonucleotide means the one that satisfies the 2 requirements, that is, it must be able to form complementary base pairing and give an -OH group serving as the origin of synthesis of complementary chain at the 3'-terminal. Accordingly, its backbone is not necessarily limited to the one via phosphodiester linkages. For example, it may be composed of a phosphothioate derivative having S in place of P as a backbone or a peptide nucleic acid based on peptide linkages. The bases may be those capable of complementary base pairing. In the nature, there are 5 bases, that is, A, C, T, G and U, but the base can be an analogue such

as bromodeoxyuridine. The oligonucleotide used in the present invention functions preferably not only as the origin of synthesis but also as a template for synthesis of complementary chain. The term polynucleotide in the present invention includes oligonucleotides. The term "polynucleotide" is used in the case where the chain length is not limited, while the term "oligonucleotide" is used to refer to a nucleotide polymer having a relatively short chain length.

The oligonucleotide according to the present invention has such a chain length as to be capable of base pairing with a complementary chain and to maintain necessary specificity under the given environment in the various reactions of synthesizing nucleic acid described below. Specifically, it is composed of 5 to 200 base pairs, more preferably 10 to 50 base pairs. The chain length of a primer recognizing the known polymerase catalyzing the sequence-dependent nucleic acid synthetic reaction is at least about 5 bases, so the chain length of the annealing part should be longer than that. In addition, a length of 10 bases or more is desired statistically in order to expect specificity as the nucleotide sequence. On the other hand, preparation of a too long nucleotide sequence by chemical synthesis is difficult, and thus the chain length described above is exemplified as a desired range. The chain length exemplified here refers to the chain length of a part annealing to a complementary chain. As described below, the

oligonucleotide according to the present invention can anneal finally to at least 2 regions individually. Accordingly, it should be understood that the chain length exemplified here is the chain length of each region constituting the oligonucleotide.

Further, the oligonucleotide according to the present invention can be labeled with a known labeling substance. The labeling substance includes binding ligands such as digoxin and biotin, enzymes, fluorescent substances and luminescent substances, and radioisotopes. The techniques of replacing a base constituting an oligonucleotide by a fluorescent analogue are also known (WO95/05391, Proc. Natl. Acad. Sci. USA, 91, 6644-6648, 1994).

Other oligonucleotides according to the present invention can also have been bound to a solid phase. Alternatively, an arbitrary part of the oligonucleotide may be labeled with a binding ligand such as biotin, and it can be immobilized indirectly via a binding partner such as immobilized avidin. When the immobilized oligonucleotide is used as the origin of synthesis, nucleic acid as the synthetic reaction product is captured by the solid phase, thus facilitating its separation. The separated product can be detected by a nucleic acid-specific indicator or by hybridization with a labeling probe. The target nucleic acid fragments can also be recovered by digesting the product with

arbitrary restriction enzymes.

The term "template" used in the present invention means nucleic acid serving as a template for synthesizing a complementary chain. A complementary chain having a nucleotide sequence complementary to the template has a meaning as a chain corresponding to the template, but the relationship between the two is merely relative. That is, a chain synthesized as the complementary chain can function again as a template. That is, the complementary chain can become a template.

The oligonucleotide useful in the present invention is not limited to the 2 regions described above and can contain an additional region. While X2 and X1c are arranged at the 3'- and 5'-terminals respectively, an arbitrary sequence can be interposed therebetween. For example, it can be a restriction enzyme recognition site, a promoter recognized by RNA polymerase, or DNA coding for ribozyme. By using it as a restriction enzyme recognition sequence, the nucleic acid having a complementary sequence alternately linked in a single-stranded chain as the synthetic product of the present invention can be cleaved into double-stranded nucleic acids of the same length. By arranging a promoter sequence recognized by RNA polymerase, the synthetic product of the present invention serves as the template to permit further transcription into RNA. By further arranging DNA coding for ribozyme, a system where the transcriptional product is

self-cleaved is realized. These additional nucleotide sequences are those functioning after formed into a double-stranded chain. Accordingly, when the single-stranded nucleic acid according to the present invention has formed a loop, these sequences do not function. They do not function until the nucleic acid is elongated and annealed in the absence of a loop to a chain having a complementary nucleotide sequence.

When a promoter is combined with the oligonucleotide based on the present invention in such a direction as to permit transcription of the synthesized region, the reaction product based on the present invention where the same nucleotide sequence is repeated realizes a highly efficient transcriptional system. By combining this system with a suitable expression system, translation into a protein is also feasible. That is, the system can be utilized for transcription and translation into protein in bacteria or animal cells or in vitro.

The oligonucleotide of the present invention having the structure described above can be chemically synthesized. Alternatively, natural nucleic acid may be cleaved with e.g. restriction enzymes and modified so as to be composed of, or ligated into, the nucleotide sequence described above.

The basic principle of the reaction for performing synthesis by utilizing the useful oligonucleotide described above in combination with DNA polymerase having the strand



displacement activity in the reaction of synthesizing nucleic acid according to the present invention is described by reference to Figs. 5 to 6. The oligonucleotide described above (FA in Fig. 5) anneals at X2 (corresponding to F2) to nucleic acid as a template, to provide the origin of synthesis of complementary chain. In Fig. 5, a complementary chain synthesized from FA as the origin of synthesis is displaced by synthesis of complementary chain (described below) from an outer primer (F3), to form a single-stranded chain (Fig. 5-A). When synthesis of complementary chain to the resulting complementary chain is further conducted, the 3'-terminal of nucleic acid synthesized as complementary chain in Fig. 5-A has a nucleotide sequence complementary to the oligonucleotide of the present invention. That is, because the 5'-terminal of the oligonucleotide of the present invention has the same sequence as a region X1c (corresponding to F1c), the 3'-terminal of the nucleic acid thus synthesized has a complementary sequence X1 (F1). Fig. 5 shows that the complementary chain synthesized from R1 as the origin of synthesis is displaced by synthesis of complementary chain by primer R3 as the origin of synthesis. Once the 3'-terminal portion is made ready for base pairing by this displacement, X1 (F1) at the 3'-terminal anneals to X1c (F1c) in the same chain, and elongation reaction with itself as a template proceeds (Fig. 5-B). Then, X2c (F2c) located at the 3'-terminal thereof is left as a loop not involving base

pairing. X2 (F2) in the oligonucleotide according to the present invention anneals to this loop, and a complementary chain is synthesized with said oligonucleotide as the origin of synthesis (Fig. 5-B). A product of complementary chain synthetic reaction with the previously synthesized product as a template is displaced by the strand displacement reaction so that it is made ready for base pairing.

By the basic constitution using one kind of oligonucleotide according to the present invention and an arbitrary reverse primer capable of conducting nucleic acid synthesis where a complementary chain synthesized with said oligonucleotide as a primer is used as a template, a plurality of nucleic acid synthetic products as shown in Fig. 6 can be obtained. As can be seen from Fig. 6, (D) is the desired nucleic acid product of the invention having complementary nucleotide sequence alternately linked in a single-stranded chain. Once converted into a single-stranded chain by treatment such as heat denaturation, the other product (E) serves again as a template for forming (D). If the product (D) as nucleic acid in the form of a double-stranded chain is converted into a single-stranded chain by heat denaturation, annealing occurs within the same chain at high probability without forming the original double-stranded chain. This is because a complementary chain having the same melting temperature ( $T_m$ ) undergoes intramolecular reaction preferentially over intermolecular

reaction. Each single-stranded chain derived from the product (D) annealed in the same chain is annealed in the same chain and returned to the state of (B), and each chain further gives one molecule of (D) and (E) respectively. By repeating these steps, it is possible to successively synthesize the nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain. The template and the product formed in 1 cycle are increased exponentially, thus making the reaction very efficient.

To realize the state of Fig. 5(A), the initially synthesized complementary chain should, in at least the portion to which the reverse primer anneals, should be ready for base pairing. This step can be achieved by an arbitrary method. That is, an outer primer (F3), which anneals to the first template at a region F3c at the 3'-side of the region F2c to which the oligonucleotide of the present invention anneals, is separately prepared. If this outer primer is used as the origin synthesis to synthesize a complementary chain by a polymerase catalyzing the strand displacement-type synthesis of complementary chain, the complementary chain synthesized from the F2c as the origin of synthesis in the invention is displaced, and as a result the region R1c to be annealed by R1 is made ready for base pairing (Fig. 5). By utilization of the strand displacement reaction, the reaction up to now can proceed under isothermal conditions.

When an outer primer is used, synthesis from the outer primer (F3) should be initiated after synthesis from F2c. In the most simple method, the concentration of the inner primer is made higher than the concentration of the outer primer. Specifically, the primers are used at usually 2- to 50-fold, preferably 4- to 10-fold different concentrations, whereby the reaction can proceed as expected. Further, the melting temperature ( $T_m$ ) of the outer primer is set to be lower than the  $T_m$  of the X1 (corresponding to F1 and R1) in the inner primer whereby the timing of synthesis can be controlled. That is, (outer primer F3 : F3c)  $\leq$  F2c/F2  $\leq$  (F1c/F1) or (outer primer/region at the 3'-side in the template)  $\leq$  (X2c : X2)  $\leq$  (X1c : X1). Here, the reason for (F2c/F2)  $\leq$  (F1c/F1) is for annealing between F1c/F1 prior to annealing of F2 to the loop. The annealing between F1c/F1 is an intramolecular reaction and may thus proceed preferentially at high probability. However, it is meaningful to consider  $T_m$  in order to give more desired reaction conditions. As a matter of course, similar conditions should be considered even in the design of a reverse primer. By using such a relationship, statistically ideal reaction conditions can be achieved. If other conditions are fixed, melting temperature ( $T_m$ ) can be theoretically calculated by a combination of the length of an annealing complementary chain and bases constituting base-pairing. Accordingly, those skilled in the art can derive preferable conditions on the basis

of the disclosure of this specification.

Further, the phenomenon called contiguous stacking can also be applied for controlling timing of annealing of the outer primer. Contiguous stacking is a phenomenon in which an oligonucleotide not capable of annealing independently is made capable of annealing upon being contiguous to the part of a double-stranded chain (Chiara Borghesi-Nicoletti et al., Bio Techniques, 12, 474-477 (1992)). That is, the outer primer is designed so as to be contiguous to F2c (X2c) and not to be able to anneal independently. By doing so, annealing of the outer primer does not occur until F2c (X2c) anneals, and thus the annealing of F2c (X2c) occurs preferentially. On the basis of this principle, the Examples show setting of the nucleotide sequence of an oligonucleotide necessary as a primer for a series of reactions. This step can also be achieved by denaturation under heating or with a DNA helicase.

If the template nucleic acid having F2c (X2c) is RNA, the state of Fig. 5-(A) can also be realized by a different method. For example, if this RNA chain is decomposed, R1 is made ready for base pairing. That is, F2 is annealed to F2c in RNA and a complementary chain is synthesized as DNA by a reverse transcriptase. The RNA serving as a template is decomposed by alkali denaturation or by enzymatic treatment with a ribonuclease acting on RNA in a double-stranded chain of DNA/RNA whereby the DNA synthesized from F2 is formed into a

single-stranded chain. For the enzyme selectively decomposing RNA in a double-stranded chain of DNA/RNA, the ribonuclease activity of RNase H or some reverse transcriptases can be utilized. In this manner, the reverse primer can be annealed to R1c made capable of base pairing. Accordingly, the outer primer for rendering R1c ready for base pairing becomes unnecessary.

Alternatively, the strand displacement activity of reverse transcriptase can be utilized for the strand displacement by an outer primer as described above. In this case, a reaction system can be constituted by a reverse transcriptase only. That is, using RNA as a template, it is made possible by a reverse transcriptase to synthesize a complementary chain from F2 annealing to F2c in the template and to synthesize a complementary chain from the outer primer F3 as the origin of synthesis annealing to F3c located at the 3'-side of F2c and to simultaneously displace the previously synthesized complementary chain. When the reverse transcriptase performs the reaction of synthesizing a complementary chain with DNA as the template, all the reactions of synthesizing complementary chains including the synthesis of a complementary chain with R1 as the origin of synthesis annealing to R1c in the displaced complementary chain as the template, the synthesis of a complementary chain with R3 as the origin of synthesis annealing to R3c located at the 3'-side of

R1c and the simultaneous displacement reaction, proceed by the reverse transcriptase. If it is not possible to expect that the reverse transcriptase exhibits the DNA/RNA strand displacement activity under given reaction conditions, a DNA polymerase having the strand displacement activity described above may be combined. The mode of obtaining a first single-stranded nucleic acid with RNA as a template as described above constitutes a preferable mode of the present invention. On the other hand, if a DNA polymerase such as Bca DNA polymerase having both strand displacement activity and reverse transcriptase activity is used, not only synthesis of a first single-stranded nucleic acid from RNA but also subsequent reaction with DNA as a template can proceed similarly by the same enzyme.

The reaction system described above brings about various variations inherent in the present invention by utilization of the reverse primer having a specific structure. The most effective variation is described below. That is, the oligonucleotide constituted as described in [5] is used as the reverse primer in the most advantageous mode of the present invention. The oligonucleotide in [5] is an oligonucleotide wherein arbitrary regions R2c and R1c in a complementary chain synthesized with F2 as a primer are X2c and X1c respectively. By use of such a reverse primer, a series of reactions for forming a loop and for synthesizing and displacing a complementary chain

from this loop occur in both the sense and antisense chains (forward side and reverse side). As a result, the reaction efficiency for synthesis of the nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain according to the present invention is greatly improved while a series of these reactions are feasible under isothermal conditions. Hereinafter, this mode is described in more detail by reference to Figs. 1 to 3 where this mode is summarized.

In the following mode, 2 kinds of oligonucleotides based on the present invention are prepared. For explanation, these are designated FA and RA. The regions constituting FA and RA are as follows:

	X2	X1c
FA	F2	F1c
RA	R2	R1c

Here, F2 is a complementary nucleotide sequence to a region F2c in nucleic acid as the template. R2 is a nucleotide sequence complementary to an arbitrary region R2c contained in a complementary chain synthesized with F2 as a primer. F1c and R1c are arbitrary nucleotide sequences located downward from F2c and R2c respectively. The distance between F2 and R2 may be arbitrary. Even if its length is about 1 kbp, sufficient synthesis is feasible under suitable conditions, though depending on the synthetic ability of DNA polymerase to perform



the synthesis of a complementary chain. Specifically, when Bst DNA polymerase is used, the desired product is certainly synthesized if the distance between F2 and R2c is 800 bp, preferably 500 bp or less. In PCR involving temperature cycle, the reduction in the enzyme activity by the stress of temperature change is considered to reduce the efficiency of synthesis of a long nucleotide sequence. In a preferable mode of the present invention, the temperature cycle in the step of amplifying nucleic acid is not required, and thus the synthesis and amplification of an even long nucleotide sequence can be certainly achieved.

First, F2 in FA is annealed to nucleic acid as a template and used as the origin of synthesis of a complementary chain. The subsequent reaction steps until Fig. 1(4) are the same as in the previously described basic mode (Fig. 5) in the present invention. The sequence annealed as F3 in Fig. 1(2) is the outer primer described above. A DNA polymerase for conducting the strand displacement-type synthesis of a complementary chain with this primer as the origin of synthesis is used so that the complementary chain synthesized from FA is displaced and made ready for base pairing.

When R2c is made ready for base pairing in (4), RA as a reverse primer anneals thereto in the combination of R2c/R2. Synthesis of a complementary chain with this site as the origin of synthesis proceeds until the chain reaches Flc at the

5'-terminal of FA. Following this reaction of synthesizing a complementary chain, the outer primer R3 for displacement anneals thereto to synthesize a complementary chain, during which strand displacement also proceeds so that the complementary chain synthesized from RA as the origin of synthesis is displaced. In the complementary chain thus displaced, RA is located at the 5'-side thereof and a sequence complementary to FA is located at the 3'-terminal thereof.

At the 3'-side of the single-stranded nucleic acid thus displaced, there is a sequence F1 complementary to F1c in the same chain. F1 rapidly anneals to F1c in the same molecule to initiate synthesis of a complementary chain. When the 3'-terminal (F1) anneals to F1c in the same chain, a loop containing F2c is formed. As is also evident from Fig. 2-(7), the part of this loop remains ready for base pairing. The oligonucleotide FA of the invention having a nucleotide sequence complementary to F2c anneals to the part of this loop and acts as the origin of synthesis of a complementary chain (7). Synthesis of a complementary chain from the loop proceeds while the reaction product in the previously initiated complementary chain synthesis from F1 is displaced. As a result, the complementary chain synthesized with itself as the template is made ready for base pairing again at the 3'-terminal. This 3'-terminal is provided with a region R1 capable of annealing to R1c in the same chain, and the two are annealed preferentially

due to the rapid intramolecular reaction. The same reaction as the above-described reaction starting from the 3'-terminal synthesized with FA as a template proceeds in this region as well. As a result, the nucleic acid having complementary nucleotide sequences linked alternately in the same single-stranded chain according to the present invention is continued to be extended from R1 as the starting point at the 3'-terminal by successive synthesis of a complementary chain and subsequent displacement thereof. Because R2c is always contained in the loop formed by intramolecular annealing of the 3'-terminal R1, the oligonucleotide (RA) provided with R2 anneals to the loop at the 3'-terminal in the subsequent reaction.

When attention is paid to nucleic acid synthesized as complementary chain from the oligonucleotide annealing to the loop in the single-stranded nucleic acid elongated with itself as the template, synthesis of the nucleic acid having complementary nucleotide sequences linked alternately in the same single-stranded chain according to the present invention also proceeds here. That is, synthesis of a complementary chain from the loop is completed when it reached RA in e.g. Fig. 2-(7). Then, when the nucleic acid displaced by this nucleic acid synthesis initiates synthesis of complementary chain (Fig. 3-(8)), the reaction reaches the loop which was once the origin of synthesis, and displacement is initiated again. In this manner, the nucleic acid initiated to be synthesized from the

loop is also displaced, and as a result, the 3'-terminal R1 capable of annealing in the same chain is obtained (Fig. 3-(10)). This 3'-terminal R1 anneals to R1c in the same chain to initiate synthesis of complementary chain. This reaction is the same as in Fig. 2-(7) except that F is used in place of R. Accordingly, the structure shown in Fig. 3-(10) can function as a new nucleic acid which continues self-elongation and new nucleic acid formation.

The reaction of synthesizing nucleic acid, initiated from the nucleic acid shown in Fig. 3-(10), causes elongation from the 3'-terminal F1 as the origin of synthesis, as opposed to the reaction described above. That is, in the present invention, as one nucleic acid is elongated, the reaction of continuing to supply a new nucleic acid initiating elongation separately proceeds. Further, as the chain is elongated, a plurality of loop-forming sequences are brought about not only at the terminal but also in the same chain. When these loop-forming sequences are made ready for base pairing by the strand displacement synthetic reaction, an oligonucleotide anneals thereto to serve as a base for the reaction of forming a new nucleic acid. Further efficient amplification is achieved by the synthetic reaction starting not only at the terminal but also in the chain. The oligonucleotide RA based on the present invention is combined as the reverse primer as described above whereby elongation and subsequent formation of a new nucleic

acid occur. Further, in the present invention, this newly formed nucleic acid itself is elongated and brings about subsequent formation of a new nucleic acid. A series of these reactions continue theoretically permanently to achieve very efficient amplification of nucleic acid. In addition, the reaction in the present invention can be conducted under isothermal conditions.

The reaction products thus accumulated possess a structure having a nucleotide sequence between F1 and R1 and its complementary sequence linked alternately therein. However, both the terminals of the repeating unit have a region consisting of the successive nucleotide sequences F2-F1 (F2c-F1c) and R2-R1 (R2c-R1c). For example, in Fig. 3-(9), the sequences (R2-F2c)-(F1-R2c)-(R1-F1c)-(F2-R2c) are linked in this order from the 5'-side. This is because the amplification reaction based on the present invention proceeds on the principle that the reaction is initiated from F2 (or R2) with an oligonucleotide as the origin of synthesis and then a complementary chain is elongated by the synthetic reaction from F1 (or R1) with the 3'-terminal as the origin of synthesis.

Here, in the most preferable mode, oligonucleotides FA and RA according to the present invention were used as oligonucleotides annealing to the part of a loop. However, even if these oligonucleotide having a limited structure are not used, the amplification reaction according to the present invention

can be carried out by use of an oligonucleotide capable of initiating the synthesis of a complementary chain from the loop. That is, the elongating 3'-terminal, once displaced by a complementary chain synthesized from the loop, gives the part of a loop again. Because the nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain is always used as a template in the complementary chain synthesis starting from the loop, it is evident that the nucleic acid desired in the present invention can be synthesized. However, the nucleic acid thus synthesized performs synthesis of a complementary chain by forming a loop after displacement, but there is no 3'-terminal available for subsequent formation of a loop, and thus it cannot function as a new template. Accordingly, the product in this case, unlike nucleic acid initiated to be synthesized by FA or RA, cannot be expected to be exponentially amplified. From this reason, an oligonucleotide having the structure of FA or RA is useful for highly efficient synthesis of nucleic acid based on the present invention.

A series of these reactions proceed by adding the following components to single-stranded nucleic acid as a template and then incubating the mixture at such a temperature that the nucleotide sequence constituting FA and RA can form stable base pairing with its complementary nucleotide sequence while the enzyme activity can be maintained.

- 4 kinds of oligonucleotides:

FA,

RA,

outer primer F3, and

outer primer R3,

- DNA polymerase for performing the strand displacement-type synthesis of complementary chain,
- an oligonucleotide serving as a substrate for DNA polymerase.

Accordingly, temperature cycle such as in PCR is not necessary. The stable base pairing referred to herein means a state in which at least a part of an oligonucleotide present in the reaction system can give the origin of synthesis of complementary chain. For example, the desired condition for bringing about stable base pairing is to set lower than melting temperature ( $T_m$ ). Generally, melting temperature ( $T_m$ ) is regarded as the temperature at which 50 % of nucleic acids having mutually complementary nucleotide sequences are base-paired. Setting at melting temperature ( $T_m$ ) or less is not an essential condition in the present invention, but is one of the reaction conditions to be considered for attaining high efficiency of synthesis. If nucleic acid to be used as a template is a double-stranded chain, the nucleic acid should, in at least a region to which the oligonucleotide anneals, be made ready for base pairing. For this, heat denaturation is generally conducted, and this may be conducted only once as pretreatment

before the reaction is initiated.

This reaction is conducted in the presence of a buffer giving suitable pH to the enzyme reaction, salts necessary for annealing or for maintaining the catalytic activity of the enzyme, a protective agent for the enzyme, and as necessary a regulator for melting temperature ( $T_m$ ). As the buffer, e.g. Tris-HCl having a buffering action in a neutral to weakly alkaline range is used. The pH is adjusted depending on the DNA polymerase used. As the salts, KCl, NaCl,  $(NH_4)_2SO_4$  etc. are suitably added to maintain the activity of the enzyme and to regulate the melting temperature ( $T_m$ ) of nucleic acid. The protective agent for the enzyme makes use of bovine serum albumin or sugars. Further, dimethyl sulfoxide (DMSO) or formamide is generally used as the regulator for melting temperature ( $T_m$ ). By use of the regulator for melting temperature ( $T_m$ ), annealing of the oligonucleotide can be regulated under limited temperature conditions. Further, betaine (N,N,N-trimethylglycine) or a tetraalkyl ammonium salt is also effective for improving the efficiency of strand displacement by virtue of its isostabilization. By adding betaine in an amount of 0.2 to 3.0 M, preferably 0.5 to 1.5 M to the reaction solution, its promoting action on the nucleic acid amplification of the present invention can be expected. Because these regulators for melting temperature act for lowering melting temperature, those conditions giving suitable



stringency and reactivity are empirically determined in consideration of the concentration of salts, reaction temperature etc.

An important feature in the present invention is that a series of reactions do not proceed unless the positional relationship of a plurality of regions is maintained. By this feature, unspecific synthetic reaction accompanied by unspecific synthesis of complementary chain is effectively prevented. That is, even if a certain unspecific reaction occurs, the possibility for the product to serve as a starting material in the subsequent amplification step is minimized. Further, the regulation of the progress of reactions by many regions brings about the possibility that a detection system capable of strict identification of the desired product in analogous nucleotide sequences can be arbitrarily constituted.

This feature can be utilized for detection of mutations in a gene. In the mode of the invention where the outer primer is used, 4 primers, that is, 2 outer primers and 2 primers consisting of the oligonucleotides of the present invention, are used. That is, unless the 6 regions contained in the 4 oligonucleotides work as designed, the synthetic reaction of the present invention do not proceed. In particular, the sequences of the 3'-terminal of each oligonucleotide as the origin of synthesis of complementary chain and of the 5'-terminal of the Xlc region where the complementary chain serves

as the origin of synthesis are important. Hence, these important sequences is designed so as to correspond to a mutation to be detected, and the synthetic reaction product of the present invention is observed whereby the presence or absence of a mutation such as base deletion or insertion, or genetic polymorphism such as SNPs can be comprehensively analyzed. Specifically, bases estimated to have a mutation or polymorphism are designed so as to correspond to the vicinity of the 3'-terminal of an oligonucleotide as the origin of synthesis of complementary chain, or of 5'-terminal thereof when a complementary chain is the origin of synthesis. If a mismatch is present at the 3'-terminal as the origin of synthesis of complementary chain or in its vicinity, the reaction of synthesizing a complementary chain to nucleic acid is significantly inhibited. In the present invention, a high degree of amplification reaction is not achieved unless the structure of the terminals of a product in the initial reaction brings about repeated reactions. Accordingly, even if erroneous synthesis occurs, complementary chain synthesis constituting amplification reaction is always interrupted in some of the steps, and thus a high degree of amplification reaction does not occur in the presence of a mismatch. As a result, the mismatch effectively inhibits amplification reaction, and an accurate result is finally brought about. That is, it can be said that the amplification reaction of nucleic

acid based on the present invention has a highly completed mechanism for checking the nucleotide sequence. These features are an advantage hardly expectable in e.g. the PCR method where amplification reaction is performed in mere 2 regions.

The region X1c characterizing the oligonucleotide used in the present invention can serve as the origin of synthesis after a complementary sequence is synthesized, and this complementary sequence anneals to the sequence X1 in the same newly synthesized chain whereby synthetic reaction with itself as a template proceeds. Therefore, even if the so-called primer dimer which is often problematic in the prior art is formed, this oligonucleotide does not form a loop. Accordingly, unspecific amplification attributable to the primer dimer cannot occur theoretically, and thus the present oligonucleotide contributes to an improvement in the specificity of the reaction.

Further, according to the present invention, the outer primers shown as F3 (Fig. 1-(2)) or R3 (Fig. 2-(5)) are combined whereby a series of the reactions described above can be conducted under isothermal conditions. That is, the present invention provides a method of amplifying nucleic acid having complementary sequences linked alternately in a single-stranded chain, which comprises the steps shown in item 9 above. In this method, temperature conditions where stable annealing occurs between F2c/F2, between R2c/R2, between F1c/F1, and

between R1c/R1 are selected, and preferably F3c/F3 and R3c/R3 are set up to be annealed by the phenomenon of contiguous stacking facilitated by annealing of F2c/F2 and R2c/R2, respectively.

In the present invention, the terms "synthesis" and "amplification" of nucleic acid are used. The synthesis of nucleic acid in the present invention means the elongation of nucleic acid from an oligonucleotide serving as the origin of synthesis. If not only this synthesis but also the formation of other nucleic acid and the elongation reaction of this formed nucleic acid occur continuously, a series of these reactions is comprehensively called amplification.

The single-stranded nucleic acid which is provided at the 3'-terminal thereof with a region F1 capable of annealing to a part F1c in the same chain and which upon annealing of the region F1 to F1c in the same chain, is capable of forming a loop containing a region F2c capable of base pairing is an important element of the present invention. Such a single-stranded nucleic acid can also be supplied on the following principle. That is, the synthesis of a complementary chain is allowed to proceed on the basis of a primer having the following structure. 5'-[region X1 annealing to region X1c located in primer]-[loop forming sequence ready for base pairing]-[region X1c]-[region having a sequence complementary to a template]-3'

As the region having a sequence complementary to a

template, two nucleotide sequences, that is, a nucleotide sequence (primer FA) complementary to F1 and a nucleotide sequence (primer RA) complementary to R1c, are prepared. The nucleotide sequence constituting nucleic acid to be synthesized contains a nucleotide sequence extending from the region F1 to the region R1c and a nucleotide sequence extending from the region R1 having a nucleotide sequence complementary to this nucleotide sequence to the region F1c. X1c and X1 capable of annealing in the inside of the primer can be arbitrary sequences. However, in a region between primers FA and RF, the sequence of the region X1c/X1 is made preferably different.

First, the synthesis of a complementary chain by the primer FA from the region F1 in template nucleic acid is conducted. Then, the region R1c in the synthesized complementary chain is made ready for base pairing, to which the other primer is annealed to form the origin of synthesis of complementary chain. The 3'-terminal of the complementary chain synthesized in this step has a nucleotide sequence complementary to the primer FA constituting the 5'-terminal of the initially synthesized chain, so it has been provided at the 3'-terminal thereof with the region X1 which anneals to the region X1c in the same chain to form a loop. The characteristic 3'-terminal structure according to the present invention is thus provided, and the subsequent reaction constitutes the reaction system shown previously as the most preferable mode.

The oligonucleotide annealing to the portion of the loop is provided at the 3'-terminal thereof with the region X2 complementary to the region X2c located in the loop and at the 5'-terminal thereof with the region X1. In the previous reaction system, primers FA and RA were used to synthesize a chain complementary to template nucleic acid thereby giving a loop structure to the 3'-terminal of the nucleic acid. In this method, the terminal structure characteristic of the present invention is provided by the short primers. In this mode, on the other hand, the whole of a nucleotide sequence constituting a loop is provided as a primer, and synthesis of this longer primer is necessary.

If a nucleotide sequence containing restriction enzyme recognition regions is used as a reverse primer, a different mode according to the present invention can be constituted. The reaction with a reverse primer containing a restriction enzyme recognition sequence is specifically described by reference to Fig. 6. When Fig. 6-(D) is completed, a nick is generated by a restriction enzyme corresponding to a restriction enzyme recognition site in the reverse primer. The strand displacement-type reaction of synthesizing complementary chain is initiated from this nick as the origin of synthesis. Because the reverse primers are located at both the terminals of a double-stranded nucleic acid constituting (D), the reaction of synthesizing complementary chain is also initiated from both

the terminals. Though basically based on the SDA method described as the prior art, the nucleotide sequence serving as a template has a structure having complementary nucleotide sequences alternately linked according to the present invention so that the nucleic acid synthetic system unique to the present invention is constituted. A part serving as a complementary chain of the reverse primer to be nicked should be designed to incorporate a dNTP derivative such that it is rendered nuclease resistance to prevent cleavage of the double-stranded chain by the restriction enzyme.

It is also possible to insert a promoter for RNA polymerase into the reverse primer. Transcription from both the terminals in Fig. 6-(D) is performed by a RNA polymerase recognizing this promoter in this case too similar to the previous mode where the SDA method was applied.

The nucleic acid synthesized in the present invention is a single-stranded chain but is composed of partial complementary sequences, and thus the majority of these sequences are base-paired. By use of this feature, the synthesized product can be detected. By carrying out the method of synthesizing nucleic acid according to the present invention in the presence of a fluorescent pigment as a double-stranded chain-specific intercalater such as ethidium bromide, SYBR Green I or Pico Green, the increased density of fluorescence is observed as the product is increased. By monitoring it, it

is possible to trace the real-time synthetic reaction in a closed system. Application of this type of detection system to the PCR method is also considered, but it is deemed that there are many problems because the signal from the product cannot be distinguished from signals from primer dimers etc. However, when this system is applied to this invention, the possibility of increasing unspecific base pairing is very low, and thus high sensitivity and low noises can be simultaneously expected. Similar to use of the double-stranded chain-specific intercalater, the transfer of fluorescent energy can be utilized for a method of realizing a detection system in a uniform system.

The method of synthesizing nucleic acid according to the present invention is supported by the DNA polymerase catalyzing the strand displacement-type reaction for synthesis of complementary chain. During the reaction described above, a reaction step not necessarily requiring the strand displacement-type polymerase is also contained. However, for simplification of a constitutional reagent and in an economical viewpoint, it is advantageous to use one kind of DNA polymerase. As this kind of DNA polymerase, the following enzymes are known. Further, various mutants of these enzymes can be utilized in the present invention insofar as they have both the sequence-dependent activity for synthesis of complementary chain and the strand displacement activity. The mutants



referred to herein include those having only a structure bringing about the catalytic activity required of the enzyme or those with modifications to catalytic activity, stability or thermostability by e.g. mutations in amino acids.

Bst DNA polymerase

Bca (exo-)DNA polymerase

DNA polymerase I Klenow fragment

Vent DNA polymerase

Vent (exo-)DNA polymerase (Vent DNA polymerase deficient in exonuclease activity)

Deep Vent DNA polymerase

Deep Vent(exo-)DNA polymerase (Deep Vent DNA polymerase deficient in exonuclease activity)

$\Phi$ 29 phage DNA polymerase

MS-2 phage DNA polymerase

Z-Taq DNA polymerase (Takara Shuzo Co., Ltd.)

KOD DNA polymerase (Toyobo Co., Ltd.)

Among these enzymes, Bst DNA polymerase and Bca (exo-) DNA polymerase are particularly desired enzymes because they have a certain degree of thermostability and high catalytic activity. The reaction of this invention can be carried isothermally in a preferred embodiment, but because of the adjustment of melting temperature ( $T_m$ ) etc., it is not always possible to utilize temperature conditions desired for the stability of the enzyme. Accordingly, it is one of the desired

conditions that the enzyme is thermostable. Although the isothermal reaction is feasible, heat denaturation may be conducted to provide nucleic acid as a first template, and in this respect too, utilization of a thermostable enzyme broadens selection of assay protocol.

Vent (exo-) DNA polymerase is an enzyme having both strand displacement activity and a high degree of thermostability. It is known that the complementary chain synthetic reaction involving strand displacement by DNA polymerase is promoted by adding a single strand binding protein (Paul M. Lizardi et al., Nature Genetics, 19, 225-232, July, 1998). This action is applied to the present invention, and by adding the single strand binding protein, the effect of promoting the synthesis of complementary chain can be expected. For example, T4 gene 32 is effective as a single strand binding protein for Vent (exo-) DNA polymerase.

For DNA polymerase free of 3'-5' exonuclease activity, there is a known phenomenon where the synthesis of complementary chain does not stop at the 5'-terminal of a template, resulting in generation of a one-base protrusion. In the present invention, this phenomenon is not preferable because when synthesis of the complementary chain reaches the terminal, the sequence of the 3'-terminal leads to initiation of next synthesis of complementary chain. However, because a base "A" is added at high probability to the 3'-terminal by the

DNA polymerase, the sequence may be selected such that the synthesis from the 3'-terminal starts at "A", so that there is no problem if an additional base is added erroneously by dATP. Further, even if the 3'-terminal is protruded during synthesis of complementary chain, the 3'→5' exonuclease activity can be utilized for digesting the protrusion to make it blunt-ended. For example, since natural Vent DNA polymerase has this activity, this enzyme may be used as a mixture with Vent (exo-) DNA polymerase in order to solve this problem.

Various reagents necessary for the method of synthesizing or amplifying nucleic acid according to the present invention may be previously packaged and provided as a kit. Specifically, a kit is provided for the present invention, comprising various kinds of oligonucleotides necessary as primers for synthesis of complementary chain and as outer primers for displacement, dNTP as a substrate for synthesis of complementary chain, a DNA polymerase for carrying out the strand displacement-type synthesis of complementary chain, a buffer giving suitable conditions to the enzyme reaction, and as necessary reagents necessary for detection of synthetic reaction products. In particular, the addition of reagents is necessary during the reaction in a preferable mode of the present invention, and thus the reagents necessary for one reaction are supplied after pipetted into reaction vessel, whereby the reaction can be initiated by adding only a sample. By constituting a system

in which the reaction product can be detected in situ in a reaction vessel by utilizing a luminescent signal or a fluorescent signal, it is not necessary to open and shut the vessel after reaction. This is very desirable for prevention of contamination.

The nucleic acid having complementary nucleotide sequences alternately linked in a single-stranded chain, synthesized according to the present invention, has e.g. the following usefulness. The first feature is use of an advantage resulting from the special structure having complementary sequences in one molecule. This feature can be expected to facilitate detection. That is, there is a known system for detecting nucleic acid wherein its signal is varied depending on base pairing with a complementary nucleotide sequence. For example, by combination with the method of utilizing a double-stranded chain-specific intercalater as a detector as described above, a detection system making full use of the characteristics of the synthetic product of the present invention can be realized. If the synthetic reaction product of the present invention is once heat-denatured in said detection system and returned to the original temperature, intramolecular annealing occurs preferentially thus permitting complementary sequences to be rapidly base-paired. If there are unspecific reaction products, they have not complementary sequences in the molecule so that after separated by heat

denaturation into 2 or more molecules, they cannot immediately be returned to the original double-stranded chain. By providing the step of heat denaturation before detection, noises accompanying the unspecific reaction can be reduced. If the DNA polymerase not resistant to heat is used, the step of heat denaturation has the meaning of termination of the reaction and is thus advantageous for the control of reaction temperature.

The second feature is to always form a loop capable of base pairing. The structure of a loop capable of base pairing is shown in Fig. 4. As can be seen from Fig. 4, the loop is composed of the nucleotide sequence F2c (X2c) which can be annealed by the primer and a nucleotide sequence intervening between F2c-F1c (X1c). The sequence between F2c-F1c (or between X2c-X1c in a more universal form) is a nucleotide derived sequence derived from the template. Accordingly, if a probe having a complementary nucleotide sequence is hybridized with this region, template-specific detection is feasible. In addition, this region is always ready for base pairing, and therefore, heat denaturation prior to hybridization is not necessary. The nucleotide sequence constituting a loop in the amplification reaction product in the present invention may have an arbitrary length. Accordingly, if hybridization with a probe is desired, a region to be annealed by the primer and a region to be hybridized by the probe are arranged separately

to prevent their competition, whereby ideal reaction conditions can be constituted.

According to a preferable mode of the present invention, a large number of loops capable of base pairing are given in a single strand of nucleic acid. This means that a large number of probes can be hybridized with one molecule of nucleic acid to permit highly sensitive detection. It is thus possible to realize not only the improvement of sensitivity but also a method of detecting nucleic acid based on a special reaction principle such as aggregation. For example, a probe immobilized onto fine particles such as polystyrene latex is added to the reaction product of the present invention, the aggregation of latex particles is observed as the hybridization of the product with the probe proceeds. Highly sensitive and quantitative observation is feasible by optically measuring the strength of the aggregation. Because the aggregation can also be observed with the naked eyes, a reaction system not using an optical measuring device can also be constituted.

Further, the reaction product of the present invention permitting many labels to be bound thereto per nucleic acid molecule enables chromatographic detection. In the field of immunoassay, an analytical method (immuno chromatography) using a chromatographic medium utilizing a visually detectable label is used practically. This method is based on the principle that an analyte is sandwiched between an antibody immobilized on a

chromatographic medium and a labeled antibody, and the unreacted labeled component is washed away. The reaction product of the present invention makes this principle applicable to analysis of nucleic acid. That is, a labeled probe toward the part of a loop is prepared and immobilized onto a chromatographic medium to prepare a capturing probe for trapping thereby permitting analysis in the chromatographic medium. As the capturing probe, a sequence complementary to the part of the loop can be utilized. Since the reaction product of the present invention has a large number of loops, the product binds to a large number of labeled probes to give a visually recognizable signal.

The reaction product according to the present invention always giving a region as a loop capable of base pairing enables a wide variety of other detection systems. For example, a detection system utilizing surface plasmon resonance using an immobilized probe for this loop portion is feasible. Further, if a probe for the loop portion is labeled with a double-stranded chain-specific intercalater, more sensitive fluorescent analysis can be conducted. Alternatively, it is also possible to positively utilize the ability of the nucleic acid synthesized by the present invention to form a loop capable of base pairing at both the 3'- and 5'-sides. For example, one loop is designed to have a common nucleotide sequence between a normal type and an abnormal type, while the other loop is

designed to generate a difference therebetween. It is possible to constitute a characteristic analytic system in which the presence of a gene is confirmed by the probe for the common portion while the presence of an abnormality is confirmed in the other region. Because the reaction of synthesizing nucleic acid according to the present invention can also proceed isothermally, it is a very important advantage that real-time analysis can be effected by a general fluorescent photometer. Heretofore, the structure of nucleic acid to be annealed in the same chain is known. However, the nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain obtained by the present invention is novel in that it contains a large number of loops capable of base pairing with other oligonucleotides.

On the other hand, a large number of loops themselves given by the reaction product according to the present invention can be used as probes. For example, in a DNA chip, probes should be accumulated at high density in a limited area. In present technology, however, the number of oligonucleotides which can be immobilized in a certain area is limited. Hence, by use of the reaction product of the present invention, a large number of probes capable of annealing can be immobilized at high density. That is, the reaction product according to the present invention may be immobilized as probes on a DNA chip. After amplification, the reaction product may be immobilized by any techniques known



in the art, or the immobilized oligonucleotide is utilized as the oligonucleotide in the amplification reaction of the present invention, resulting in generating the immobilized reaction product. By use of the probe thus immobilized, a large number of sample DNAs can be hybridized in a limited area, and as a result, high signals can be expected.

#### Brief Description of the Drawings

Fig. 1 is an illustration of a part (1) to (4) of the reaction principle in a preferable mode of the present invention.

Fig. 2 is an illustration of a part (5) to (7) of the reaction principle in a preferable mode of the present invention.

Fig. 3 is an illustration of a part (8) to (10) of the reaction principle in a preferable mode of the present invention.

Fig. 4 is an illustration of the structure of a loop formed by the single-stranded nucleic acid according to the present invention.

Fig. 5 is an illustration of a part (A) to (B) in a basic mode of the present invention.

Fig. 6 is an illustration of a part (C) to (D) in a basic mode of the present invention.

Fig. 7 is a drawing showing the positional relationship

of each nucleotide sequence constituting an oligonucleotide in the target nucleotide sequence of M13mp18.

Fig. 8 is a photograph showing the result of agarose electrophoresis of a product obtained by the method of synthesizing single-stranded nucleic acid with M13mp18 as a template according to the present invention.

Lane 1: XIV size marker

Lane 2: 1 fmol M13mp18 dsDNA

Lane 3: No target

Fig. 9 is a photograph showing the result of agarose gel electrophoresis of a restriction enzyme-digested product obtained in Example 1 by the nucleic acid synthetic reaction according to the present invention.

Lane 1: XIV size marker

Lane 2: BamHI digest of the purified product

Lane 3: PvuII digest of the purified product

Lane 4: HindIII digest of the purified product

Fig. 10 is a photograph showing the result of agarose gel electrophoresis of a product obtained by the method of synthesizing single-stranded nucleic acid according to the present invention using M13mp18 as a template in the presence of betaine. 0, 0.5, 1 and 2 indicate the concentration (M) of betaine added to the reaction solution. N indicates the negative control, and -21 indicates the concentration  $10^{-21}$  mol of template DNA.

Fig. 11 is a drawing showing the positional relationship of each nucleotide sequence constituting an oligonucleotide in a target nucleotide sequence derived from HVB.

Fig. 12 is a photograph showing the result of agarose gel electrophoresis of a product obtained by the method of synthesizing single-stranded nucleic acid according to the present invention wherein HBV-M13mp18 integrated in M13mp18 was used as a template.

Lane 1: XIV size marker

Lane 2: 1 fmol HBV-M13mp18 dsDNA

Lane 3: No target

Fig. 13 is a photograph showing the result of gel electrophoresis of an alkali-denatured product obtained by the method of synthesizing single-stranded nucleic acid according to the present invention.

Lane 1: HindIII-digested fragment from lambda-phage

Lane 2: The reaction product in Example 1.

Lane 3: The reaction product in Example 3.

Fig. 14 is a photograph showing the result of agarose gel electrophoresis of a product obtained by the method of synthesizing single-stranded nucleic acid according to the present invention wherein the concentration of M13mp18 as a target was varied. The upper and lower photographs show the result of the reaction for 1 and 3 hours respectively.

Lane 1: M13mp18 dsDNA  $1 \times 10^{-15}$  mol/tube

Lane 2: M13mp18 dsDNA  $1 \times 10^{-16}$  mol/tube  
 Lane 3: M13mp18 dsDNA  $1 \times 10^{-17}$  mol/tube  
 Lane 4: M13mp18 dsDNA  $1 \times 10^{-18}$  mol/tube  
 Lane 5: M13mp18 dsDNA  $1 \times 10^{-19}$  mol/tube  
 Lane 6: M13mp18 dsDNA  $1 \times 10^{-20}$  mol/tube  
 Lane 7: M13mp18 dsDNA  $1 \times 10^{-21}$  mol/tube  
 Lane 8: M13mp18 dsDNA  $1 \times 10^{-22}$  mol/tube  
 Lane 9: No target  
 Lane 10: XIV size marker

Fig. 15 is a drawing showing the position of a mutation and the positional relationship of each region toward a target nucleotide sequence (target). Underlined guanine is replaced by adenine in the mutant.

Fig. 16 is a photograph showing the result of agarose gel electrophoresis of a product according to the amplification reaction of the present invention.

M: 100 bp ladder (New England Biolabs)  
 N: No template (purified water)  
 WT: 1 fmol wild-type template M13mp18  
 MT: 1 fmol mutant template M13mp18FM

Fig. 17 is a drawing showing the positional relationship of each nucleotide sequence constituting an oligonucleotide in a nucleotide sequence coding for target mRNA.

Fig. 18 is a photograph showing the result of agarose electrophoresis of a product obtained by the method of

synthesizing single-stranded nucleic acid according to the present invention using mRNA as a target.

#### Best Mode for Carrying Out the Invention

##### Example 1. Amplification of a region in M13mp18

The method of synthesizing the nucleic acid having complementary chains alternately linked in a single-stranded chain according to the present invention was attempted using M13mp18 as a template. Four kinds of primers, that is, M13FA, M13RA, M13F3, and M13R3, were used in the experiment. M13F3 and M13R3 were outer primers for displacing the first nucleic acid obtained respectively with M13FA and M13RA as the origin of synthesis. Because the outer primers are primers serving as the origin of synthesis of complementary chain after synthesis with M13FA (or M13RA), these were designed to anneal to a region contiguous to M13FA (or M13RA) by use of the phenomenon of contiguous stacking. Further, the concentrations of these primers were set high such that annealing of M13FA (or M13RA) occurred preferentially.

The nucleotide sequence constituting each primer is as shown in the Sequence Listing. The structural characteristics of the primers are summarized below. Further, the positional relationship of each region toward the target nucleotide sequence (target) is shown in Fig. 7.

Primer	Region at the 5'-side/region at the 3'-side
--------	---

M13FA            The same as region F1c in complementary chain synthesized by M13FA/complementary to region F2c in M13mp18

M13RA            The same as region R1c in complementary chain synthesized by M13RA/complementary to region R2c in complementary chain synthesized by M13FA

M13F3            Complementary to F3c contiguous to the 3'-side of region F2c in M13mp18

M13R3            Complementary to R3c contiguous to the 3'-side of region F2c in complementary chain synthesized by M13FA

By such primers, nucleic acid wherein a region extending from F1c to R1c in M13mp18, and its complementary nucleotide sequence, are alternately linked via a loop-forming sequence containing F2c in a single-stranded chain, is synthesized. The composition of a reaction solution for the method of synthesizing nucleic acid by these primers according to the present invention is shown below.

Composition of the reaction solution (in 25  $\mu$ L)

20 mM Tris-HCl pH8.8

10 mM KCl

10 mM  $(\text{NH}_4)_2\text{SO}_4$

6 mM  $\text{MgSO}_4$

0.1 % Triton X-100

5% dimethyl sulfoxide (DMSO)

0.4 mM dNTP

Primers:

800 nM M13FA/SEQ ID NO:1

800 nM M13RA/SEQ ID NO:2

200 nM M13F3/SEQ ID NO:3

200 nM M13R3/SEQ ID NO:4

Target: M13mp18 dsDNA/SEQ ID NO:5

Reaction: The above reaction solution was heated at 95 °C for 5 minutes, and the target was denatured into a single-stranded chain. The reaction solution was transferred to ice-cold water, and 4 U of Bst DNA polymerase (NEW ENGLAND Biolabs) was added thereto and the mixture was reacted at 65 °C for 1 hour. After reaction, the reaction was terminated at 80 °C for 10 minutes and transferred again to ice-cold water.

Confirmation of the reaction: 1 µl loading buffer was added to 5 µl of the above reaction solution, and the sample was electrophoresed for 1 hour at 80 mV on 2 % agarose gel (0.5 % TBE). As a molecular-weight marker, XIV (100 bp ladder, Boehringer Mannheim) was used. The gel after electrophoresis was stained with SYBR Green I (Molecular Probes, Inc.) to confirm the nucleic acid. The results are shown in Fig. 8. The respective lanes correspond to the following samples.

1. XIV size marker.
2. 1 fmol M13mp18 dsDNA.
3. No target.

In lane 3, no band was confirmed except that the unreacted primers were stained. In lane 2 in the presence of the target,

the products were confirmed as a low size band ladder, as smeared staining at high size and as a band hardly electrophoresed in the gel. Among the low-size bands, bands in the vicinity of 290 bp and 450 bp agree with the products estimated in the synthetic reaction of this invention, that is, double-stranded chains of SEQ ID NOS:11 and 12 (corresponding to double-stranded chains formed as shown in Figs. 2-(7) and 2-(10)) and a single-stranded chain of SEQ ID NO:13 (corresponding to the long single-stranded chain in Fig. 3-(9)), and it was thus confirmed that the reaction proceeds as expected. It was estimated that the electrophoresis results of the smeared pattern at high size and the band not electrophoresed were brought about because this reaction was basically a continuous reaction to permit varying molecular weights of the reaction product and further because the product has a complicated structure having a partially single-stranded chain and a double-stranded complex.

Example 2. Confirmation of the reaction products by digestion with restriction enzymes

For the purpose of clarifying the structure of the nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain obtained in Example 1 according to the present invention, the digestion of the products with restriction enzymes was conducted. If fragments are theoretically generated by digestion thereof with restriction enzymes and simultaneously the smear pattern at



high size and the band not electrophoresed as observed in Example 1 disappear, then it can be estimated that any of these products are the nucleic acid having complementary sequences linked alternately in a single-stranded chain synthesized according to the present invention.

The reaction solution (200  $\mu$ l) from 8 tubes in Example 1 was pooled and purified by treatment with phenol and precipitation with ethanol. The resulting precipitates were recovered and dissolved again in 200  $\mu$ l TE buffer, and 10  $\mu$ l aliquot was digested at 37 °C for 2 hours with restriction enzymes BamHI, PvuII, and HindIII respectively. The digest was electrophoresed for 1 hour at 80 mV on 2 % agarose gel (0.5 %TBE). As a molecular marker, Super Ladder-Low (100 bp ladder) (Gensura Laboratories, Inc.) was used. The gel after electrophoresis was stained with SYBR Green I (Molecular Probes, Inc.) to confirm the nucleic acid. The results are shown in Fig. 9. The respective lanes correspond to the following samples.

1. XIV size marker
2. BamHI digest of the purified product
3. PvuII digest of the purified product
4. HindIII digest of the purified product

It is estimated that nucleotide sequences constituting relatively short amplification products are those of SEQ ID NOS:13, 14, 15 and 16. From these nucleotide sequences, the estimated size of each fragment digested with the restriction

enzymes is as shown in Table 1. "L" in the table indicates that its position in electrophoresis is not established because L is a fragment containing a loop (single-stranded chain).

Table 1. Restriction enzyme-digested fragments of the amplification products according to the present invention

SEQ ID NO	BamHI	PvuII	HindIII
13	177 +L	56 +L	147 +L
14	15+101 +L	-	142 +L
15	171+101 +L	56 +L	147+161 +L
16	11+101+230+L	237 +L	142+170 +L
Summary	101,177,230	56,237	142,147,161,170

(11,15; not confirmed)

Because almost all bands before digestion were changed into estimated bands, it was confirmed that the object reaction products were amplified. Further, it was also shown that there were no or less unspecific products.

Example 3. Promotion of amplification reaction by addition of betaine

An experiment for examining the effect of betaine (N,N,N-trimethylglycine, Sigma) added to the amplification reaction solution on the amplification reaction of nucleic acid was conducted. Synthesis of the nucleic acid having complementary chains alternately linked in a single-stranded chain according to the present invention was conducted using M13mp18 as a template similarly in Example 1 in the presence of betaine at various concentrations. The primers used in the experiment were identical to those used in Example 1. The amount of the template DNA was  $10^{-21}$  mol (M13mp18) and water was

used as the negative control. Betaine was added at concentrations of 0, 0.5, 1 and 2 M to the reaction solution. The composition of the reaction solution is shown below.

Composition of the reaction solution (in 25  $\mu$ L)

20 mM Tris-HCl pH8.8

4 mM  $\text{MgSO}_4$

0.4 mM dNTPs

10 mM KCl

10 mM  $(\text{NH}_4)_2\text{SO}_4$

0.1% Triton X-100

Primers:

800 nM M13FA/SEQ ID NO:1

800 nM M13RA/SEQ ID NO:2

200 nM M13F3/SEQ ID NO:3

200 nM M13R3/SEQ ID NO:4

Target: M13mp18 dsDNA/SEQ ID NO:5

The polymerase, reaction conditions, and conditions for electrophoresis after the reaction were identical to those described in Example 1.

The results are shown in Fig. 10. In the reaction in the presence of betaine at a concentration of 0.5 or 1.0 M, the amount of the amplification product was increased. Further, if its concentration was increased to 2.0 M, no amplification was observed. It was thus shown that the amplification reaction was promoted in the presence of betaine at a suitable

concentration. The estimated reason that the amount of the amplification product was decreased when the concentration of betaine was 2 M was that  $T_m$  was lowered too much.

#### Example 4. Amplification of HBV gene sequence

The method of synthesizing nucleic acid according to the present invention was attempted wherein M13mp18 dsDNA having a partial sequence of HBV gene integrated therein was used as a template. Four kinds of primers, HB65FA (SEQ ID NO:6), HB65RA (SEQ ID NO:7), HBF3 (SEQ ID NO:8) and HBR3 (SEQ ID NO:9), were used in the experiment. HBF3 and HBR3 were outer primers for displacement of the first nucleic acid obtained respectively with HB65FA and HB65RA as the origin of synthesis. Because the outer primers are primers serving as the origin of synthesis of complementary chain after synthesis with HB65FA (or HB65RA), these were designed to anneal to a region contiguous to HB65FA (or HB65RA) by use of the phenomenon of contiguous stacking. Further, the concentrations of these primers were set high such that annealing of HB65FA (or HB65RA) occurred preferentially. The target sequence (430 bp) in this example, derived from HBV integrated in M13mp18, is shown in SEQ ID NO:10.

The nucleotide sequence constituting each primer is shown in the Sequence Listing. The structural feature of each primer is summarized below. Further, the positional relationship of each region toward the target nucleotide sequence (target) is shown in Fig. 11.

Primer	Region at the 5'-side/region at the 3'-side
HB65FA	The same as region F1c in complementary chain synthesized by HB65FA/complementary to region F2c in HBV-M13mp18
HB65RA	The same as region R1c in complementary chain synthesized by HB65RA/complementary to region R2c in complementary chain synthesized by HB65FA
HBf3	Complementary to F3c contiguous to the 3'-side of region F2c in HBV-M13mp18
HBR3	Complementary to R3c contiguous to the 3'-side of region F2c in complementary chain synthesized by HB65FA

By such primers, nucleic acid wherein a region extending from F1c to R1c in M13mp18 (HBV-M13mp18) having a partial sequence of HBV gene integrated therein, and its complementary nucleotide sequence, are alternately linked via a loop-forming sequence containing F2c in a single-stranded chain, is synthesized. The reaction was conducted under the same conditions as in Example 1 except that the primers described above were used, and the reaction solution was analyzed by agarose electrophoresis. The results are shown in Fig. 12. The respective lanes correspond to the following samples.

1. XIV size marker
2. 1 fmol HBV-M13mp18 dsDNA.
3. No target

Similar to Example 1, the products were confirmed only

in the presence of the target as a low size band ladder, as smeared staining at high size and as a band hardly electrophoresed in the gel (lane 2). Among the low-size bands, bands in the vicinity of 310 bp and 480 bp agree with the products estimated in the synthetic reaction of this invention, that is, double-stranded chains of SEQ ID NOS:17 and 18, and it was thus confirmed that the reaction proceeds as expected. As described in the results in Example 1, it was estimated that the smeared pattern at high size and the band not electrophoresed were caused by the structure of the synthetic product characteristic of the present invention. From this experiment, it was confirmed that the present invention can be practiced even if a different sequence (target) is used for amplification.

#### Example 5. Confirmation of the sizes of the synthetic reaction products

To confirm the structure of the nucleic acid synthesized according to the present invention, its length was analyzed by electrophoresis under alkali-denaturing conditions. 1  $\mu$ l alkaline loading buffer was added to 5  $\mu$ l of each reaction solution in the presence of the target in Example 1 or 4 and electrophoresed at 50 mA in 0.7 % agarose gel (50 mM NaOH, 1 mM EDTA) for 14 hours. As the molecular-weight size marker, HindIII-digested lambda-phage fragments were used. The gel after electrophoresis was neutralized with 1 M Tris, pH 8 and stained with SYBR Green I (Molecular Probes, Inc.) to confirm

the nucleic acid. The results are shown in Fig. 13. The respective lanes correspond to the following samples.

1. HindIII-digested fragments from lambda-phage.
2. The reaction product in Example 1.
3. The reaction product in Example 4.

When the reaction product was electrophoresed under alkali-denaturing conditions, its size in a single-stranded state could be confirmed. It was confirmed that the sizes of the major products in both Example 1 (lane 2) and Example 4 (lane 3) were within 2 kbse. Further, it was revealed that the product according to the present invention had been extended to have a size of at least 6 kbse or more within the range capable of confirmation by this analysis. In addition, it was confirmed again that bands not electrophoresed under non-denaturing conditions in Examples 1 and 4 were separated in a denatured state into individual single-stranded chains of smaller size. Example 6. Confirmation of amplification depending on the concentration of a target in the amplification of a region in M-13mp13

The influence of a varying concentration of a target on the method of synthesizing nucleic acid according to the present invention was observed. The method of synthesizing nucleic acid according to the present invention was carried out under the same conditions as in Example 1 except that the amount of M13mp18 dsDNA as the target was 0 to 1 fmol and the reaction

time was 1 hour or 3 hours. Similar to Example 1, the sample was electrophoresed in 2 % agarose gel (0.5 % TBE) and stained with SYBR Green I (Molecular Probes, Inc.) to confirm the nucleic acid. As a molecular size marker, XIV (100 bp ladder, Boehringer Mannheim) was used. The results are shown in Fig. 14 (upper: 1-hour reaction, below: 3-hour reaction). The respective lanes correspond to the following samples:

1. M13mp18 dsDNA  $1 \times 10^{-15}$  mol/tube.
2. M13mp18 dsDNA  $1 \times 10^{-16}$  mol/tube.
3. M13mp18 dsDNA  $1 \times 10^{-17}$  mol/tube.
4. M13mp18 dsDNA  $1 \times 10^{-18}$  mol/tube.
5. M13mp18 dsDNA  $1 \times 10^{-19}$  mol/tube.
6. M13mp18 dsDNA  $1 \times 10^{-20}$  mol/tube.
7. M13mp18 dsDNA  $1 \times 10^{-21}$  mol/tube.
8. M13mp18 dsDNA  $1 \times 10^{-22}$  mol/tube.
9. No target.
10. XIV size marker.

A common band among the respective lanes appears in a lower part in the electrophoretic profile and shows the unreacted stained primers. Regardless of the reaction time, no amplification product is observed in the absence of the target. A staining pattern, depending on the concentration of the target, of the amplification product was obtained only in the presence of the target. Further, the amplification product could be confirmed at lower concentration as the reaction time was



increased.

#### Example 7. Detection of a point mutation

##### (1) Preparation of M13mp18FM (mutant)

The target DNA used was M13mp18 (wild-type) and M13mp18FM (mutant). For the construction of the mutant M13mp18FM, LA PCR™ in vitro Mutagenesis Kit (Takara Shuzo Co., Ltd.) was used to replace one nucleotide for mutation. Thereafter, the sequence was confirmed by sequencing. The sequence of the F1 region is shown below:

Wild-type: CCGGGGATCCTCTAGAGTCG (SEQ ID NO:19)

Mutant: CCGGGGATCCTCTAGAGTCA (SEQ ID NO:20)

##### (2) Design of primers

The FA primers used for the wild-type and the mutant were provided at the 5'-terminal of the F1c region thereof with different nucleotide sequences, respectively. The location of the mutation and the positional relationship of each region toward the target nucleotide sequence (target) are shown in Fig. 15.

##### (3) Amplification reaction

An experiment was conducted to examine whether template-specific amplification reaction occurs using a combination of specific primers shown below by use of M13mp18 (wild-type) and M13mp18FM (mutant) as primers.

Primer set for wild-type amplification: FAd4, RAd4, F3, R3

Primer set for mutant amplification: FAMd4, RAd4, F3, R3

The nucleotide sequence of each primer is as follows:

FAd4 : CGACTCTAGAGGATCCCCGGTTTTTGTGTGTGGAATTGTGAGCGGAT (SEQ ID NO:21)

FAMd4: TGACTCTAGAGGATCCCCGGTTTTTGTGTGTGGAATTGTGAGCGGAT (SEQ ID NO:22)

RAAd4 : CGTCGTGACTGGGAAAACCCCTTTTTGTGCGGGCCTCTTCGCTATTAC (SEQ ID NO:23)

F3: ACTTTATGCTTCCGGCTCGTA (SEQ ID NO:24)

R3: GTTGGGAAGGGCGATCG (SEQ ID NO:25)

#### (4) Detection of the point mutation in M13mp18

The composition of the reaction solution is as follows:

		Final concentration
D2W	3.75 $\mu$ L	
10X Thermo pol buffer(NEB)	2.5 $\mu$ L	20 mM Tris-HCl pH 8.8 10 mM KCl 10 mM $(\text{NH}_4)_2\text{SO}_4$ 6 mM $\text{MgSO}_4$ 0.1 % TritonX-100
2.5 mM dNTP	4 $\mu$ L	400 $\mu$ M
100 mM $\text{MgSO}_4$	0.5 $\mu$ L	
4 M Betaine	6.25 $\mu$ L	1 M
M13FAd4 primer (10 pmol/ $\mu$ L) or		
M13FAMd4 primer (10 pmol/ $\mu$ L)	2 $\mu$ L	800 nM
M13RAAd4 primer (10 pmol/ $\mu$ L)	2 $\mu$ L	800 nM
M13F3 primer (10 pmol/ $\mu$ L)	0.5 $\mu$ L	200 nM

M13R3 primer (10 pmol/ $\mu$ L)	0.5 $\mu$ L	200 nM
Total amount	22 $\mu$ L	

1 fmol (2  $\mu$ l) of the target M13mp18 or M13mp18FM was added to the reaction solution and heated at 95 °C for 5 minutes whereby the target was denatured into a single-stranded chain. The reaction solution was transferred to ice-cold water, and 1  $\mu$ l (8 U) of Bst DNA polymerase (NEW ENGLAND Biolab) was added thereto and reacted for 1 hour at 68 °C or 68.5 °C. After reaction, the reaction was terminated at 80 °C for 10 minutes, and the reaction solution was transferred again to ice-cold water.

As shown in Fig. 16, when FAd4 for wild type was used as the FA primer, effective amplification was observed only in the presence of the wild-type template. On the other hand, when FAMd4 for mutant was used as the FA primer, effective amplification was observed only in the presence of the wild-type [sic.] template.

From the results described above, it was shown that the point mutation could be detected efficiently by use of the amplification reaction of the present invention.

#### Example 8. Amplification reaction of mRNA as a target

The method of synthesizing nucleic acid according to the present invention was attempted using mRNA as the target nucleic acid. To prepare the target mRNA, prostate cancer cell line LNCaP cells (ATCC No. CRL-1740) expressing prostate specific antigen (PSA) were mixed with chronic myeloid leukemia cell line

K562 cells (ATCC No. CCL-243) as non-expressing cells at 1 : 10<sup>6</sup> to 100 : 10<sup>6</sup>, followed by extraction of the total RNA by use of an RNeasy Mini kit from Qiagen (Germany). Four kinds of primers, that is, PSAFA, PSARA, PSAF3 and PSAR3, were used in the experiment. PSAF3 and PSAR3 are outer primers for displacing the first nucleic acid obtained respectively with PSAFA and PSARA as the origin of synthesis. Further, the concentrations of these primers were set high such that annealing of PSAFA (or PSARA) occurred preferentially. The nucleotide sequences constituting the respective primers are as follows.

Primer:

PSAFA: TGTTCTGATGCAGTGGGCAGCTTTAGTCTGCGGCGGTGTTCTG (SEQ ID NO: 26)

PSARA: TGCTGGGTCGGCACAGCCTGAAGCTGACCTGAAATACCTGGCCTG (SEQ ID NO: 27)

PSAF3: TGCTTGTGGCCTCTCGTG (SEQ ID NO: 28)

PSAR3: GGGTGTGGGAAGCTGTG (SEQ ID NO: 29)

The structural features of the primers are summarized below. Further, the positional relationship of each primer toward the DNA nucleotide sequence coding for the target mRNA and recognition sites of restriction enzyme Sau3AI are shown in Fig. 17.

Primer	Region at the 5'-side/region at the 3'-side
PSAFA	The same as region Flc in complementary chain

synthesized by PSAFA/complementary to region F2c in the target nucleotide sequence

PSARA            The same as region R1c in complementary chain synthesized by PSARA/complementary to region R2c in complementary chain synthesized by PSAFA

PSAF3            Complementary to F3c contiguous to the 3'-side of region F2c in the target nucleotide sequence

PSAR3            Complementary to R3c contiguous to the 3'-side of region R2c in complementary chain synthesized by PSAFA

The composition of a reaction solution for the method of synthesizing nucleic acid according to the present invention is as follows:

Composition of the reaction solution (in 25  $\mu$ L)

20 mM Tris-HCl pH 8.8

4 mM MgSO<sub>4</sub>

0.4 mM dNTPs

10 mM KCl

10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>

0.1 % Triton X-100

0.8 M betaine

5 mM DTT

1600 nM PSAFA & PSARA primer

200 nM PSAF3 & PSAR3 primer

8 U Bst DNA polymerase

100 U Rever Tra Ace (Toyobo Co., Ltd., Japan)

5 µg total RNA

All ingredients were mixed on ice. In this experiment, mRNA (single-stranded chain) is used as a target, and thus the step of making single-stranded chain by heat denaturation is not necessary. The reaction was conducted at 65 °C for 45 minutes, and the reaction was terminated at 85 °C for 5 minutes. After reaction, 5 µl of the reaction solution was electrophoresed in 2 % agarose and detected by SYBR Green I.

The results are shown in Table 18. The respective lanes correspond to the following samples.

Lane	Bst	RT	Number of LNCaP cells/10 <sup>6</sup> K562 cells
1	-	+	0
2	-	+	10
3	+	-	0
4	+	-	10
5	+	+	0
6	+	+	1
7	+	+	10

8 Sau3AI digest of 1 µL aliquot of the reaction solution in lane 6

9 Sau3AI digest of 1 µL aliquot of the reaction solution in lane 7

10 Size maker, 100 bp ladder (New England Biolabs)

In the absence of either Bst DNA polymerase or Rever Tra Ace, no amplification product could be obtained (lanes 1 to 4).

In the presence of both the enzymes, an amplification product was detected (lanes 5 to 7) if RNA derived from LNCaP was present. RNA extracted from one LNCaP cell/one million K562 cells could be detected (lane 6). When the amplification product was digested at the restriction enzyme site Sau3AI located in the inside of the target, the product was digested into a fragment of estimated size (lanes 8 and 9).

From the results described above, it was confirmed that the desired reaction product can be obtained in the method of synthesizing nucleic acid according to the present invention even if RNA is used as a target.

#### Industrial Applicability

According to the novel oligonucleotide according to the present invention and the method of synthesizing nucleic acid by using said oligonucleotide, there is provided a method of synthesizing nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain, without requiring any complicated control of temperature. A complementary chain synthesized with the oligonucleotide as a primer based on the present invention serves as the origin of synthesizing a new complementary chain with the 3'-terminal of said synthesized chain as a template. This is accompanied by formation of a loop causing annealing of a new primer, and a product of the reaction of synthesizing complementary chain

with the previously synthesized chain as a template is displaced again by synthesis of complementary chain from the loop and made ready for base pairing. The thus obtained nucleic acid synthesized with itself as a template is combined with e.g. a known nucleic acid synthesizing method such as SDA, to contribute the improvement of efficiency of nucleic acid synthesis.

According to an additional preferred mode of the present invention, there is provided a novel method of synthesizing nucleic acid, which achieves the improvement of efficiency of the known method of synthesizing nucleic acid, does not require complicated control of temperature, can be expected to attain high efficiency of amplification and can achieve high specificity. That is, the oligonucleotide based on the present invention is applied to a template chain and its complementary chain whereby nucleic acid having complementary sequences linked alternately in a single-stranded chain can be successively synthesized. This reaction continues theoretically until the starting materials necessary for synthesis are exhausted, during which new nucleic acid initiated to be synthesized from the loop continues to be formed. The elongation from the oligonucleotide having annealed to the loop performs strand displacement for supplying 3'-OH for elongation of long single-stranded nucleic acid (that is, nucleic acid having plural pairs of complementary chains linked



therein). On the other hand, the 3'-OH of the long single-stranded chain performs the reaction of synthesizing complementary chain with itself as a template whereby its elongation is achieved, during which a new complementary chain whose synthesis is initiated from the loop is displaced. Such an amplification reaction step proceeds under isothermal conditions while maintaining high specificity.

The oligonucleotides in the present invention can, when two contiguous regions are arranged as designed, function as primers for the reaction of synthesizing nucleic acid according to the present invention. This contributes significantly to the preservation of specificity. By comparison with e.g. PCR where unspecific amplification reaction is initiated by unspecific misannealing regardless of the intended positional relationship of 2 primers, it can be easily explained that high specificity can be expected in the present invention. This feature can be utilized to detect SNPs highly sensitively and accurately.

The characterizing feature of the present invention lies in that such reaction can be easily achieved by a very simple constitution of reagents. For example, the oligonucleotide according to the present invention has a special structure, but this is a matter of selection of nucleotide sequence, and it is a simple oligonucleotide as substance. Further, in a preferred mode, the reaction can proceed by only a DNA polymerase

catalyzing the strand displacement-type reaction of synthesizing complementary chain. Further, if the present invention is carried out with RNA as a template, a DNA polymerase such as Bca DNA polymerase having both reverse transcriptase activity and strand displacement-type DNA polymerase activity is used so that all enzyme reactions can be conducted by the single enzyme. The reaction principle of realizing a high degree of nucleic acid amplification reaction by such simple enzyme reaction is not known. Even for the application of the present invention to a known nucleic acid synthesizing reaction such as SDA, no additional enzyme is necessary for their combination, and such a simple combination with the oligonucleotide based on the present invention can be applied to various reaction systems. Accordingly, it can be said that the method of synthesizing nucleic acid according to the present invention is also advantageous in respect of cost.

As described above, the method of synthesizing nucleic acid according to the present invention and the oligonucleotide therefor provide a new principle of simultaneously solving a plurality of difficult problems such as operativeness (temperature control is not necessary), improvement of efficiency of synthesis, economization, and high specificity.

## CLAIMS

1. A method of synthesizing nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain, comprising:

- a) the step of giving nucleic acid which is provided at the 3'-terminal thereof with a region F1 capable of annealing to a part F1c in the same chain and which upon annealing of the region F1 to F1c, is capable of forming a loop containing a region F2c capable of base pairing,
- b) the step of performing synthesis of a complementary chain wherein the 3'-terminal of F1 having annealed to F1c serves as the origin of synthesis,
- c) the step of annealing, to a region F2c, of an oligonucleotide provided with the 3'-terminal thereof with F2 consisting of a sequence complementary to the region F2c, followed by synthesis, with said oligonucleotide as the origin of synthesis, of a complementary chain by a polymerase catalyzing the strand displacement reaction of synthesizing a complementary chain to displace the complementary chain synthesized in step b), and
- d) the step of annealing, to the complementary chain displaced in step c) to be ready for base pairing, of a polynucleotide provided at the 3'-terminal thereof with a sequence complementary to an arbitrary region in said chain synthesized in step c), followed by synthesis, with said 3'-terminal as the origin of synthesis, of a complementary chain by a polymerase

catalyzing the strand displacement reaction of synthesizing a complementary chain to displace the complementary chain synthesized in step c).

2. The method according to claim 1, wherein in step d), the origin of synthesis is a region R1 present at the 3'-terminal in the same chain and capable of annealing to a region R1c, and a loop containing the region R2c capable of base pairing is formed by annealing R1 to R1c.

3. An oligonucleotide composed of at least two regions X2 and X1c below, and X1c is linked to the 5'-side of X2,

X2: a region having a nucleotide sequence complementary to an arbitrary region X2c in nucleic acid having a specific nucleotide sequence, and

X1c: a region having substantially the same nucleotide sequence as in a region X1c located at the 5'-side of the region X2c in nucleic acid having a specific nucleotide sequence.

4. The method according to claim 1, wherein the nucleic acid in step a) is second nucleic acid provided by the following steps:

i) the step of annealing, to a region F2c in nucleic acid serving as a template, of a region F2 in the oligonucleotide described in claim 3 wherein the region X2 is a region F2 and the region X1c is a region F1c,

ii) the step of synthesizing first nucleic acid having a nucleotide sequence complementary to the template wherein F2

in the oligonucleotide serves as the origin of synthesis,  
iii) the step of rendering an arbitrary region in the first  
nucleic acid synthesized in step ii) ready for base pairing,  
and

iv) the step of annealing an oligonucleotide having a nucleotide  
sequence complementary to the region made ready for base pairing  
in the first nucleic acid in step iii), followed by synthesizing  
second nucleic acid with said oligonucleotide as the origin of  
synthesis and rendering F1 at the 3'-terminal thereof ready for  
base pairing.

5. The method according to claim 4, wherein the region enabling  
base pairing in step iii) is R2c, and the oligonucleotide in  
step iv) is the oligonucleotide described in claim 3 wherein  
the region X2c is a region R2c and the region X1c is a region  
R1c.

6. The method according to claim 4 or 5, wherein the step of  
rendering base pairing ready in steps iii) and iv) is conducted  
by the strand displacement synthesis of complementary chain by  
a polymerase catalyzing the strand displacement reaction of  
synthesizing complementary chain wherein an outer primer  
annealing to the 3'-side of F2c in the template and an outer  
primer annealing to the 3'-side of the region used as the origin  
of synthesis in step iv) for the first nucleic acid serve as  
the origin of synthesis.

7. The method according to claim 6, wherein the melting

temperature of each oligonucleotide and its complementary region in the template used in the reaction is in the following relationship under the same stringency:

(outer primer/region at the 3'-side in the template)  $\leq$  (F2c/F2 and R2c/R2)  $\leq$  (F1c/F1 and R1c/R1).

8. The method according to any one of claims 4 to 7, wherein the nucleic acid serving as the template is RNA, and the synthesis of complementary chain in step ii) is conducted by an enzyme having a reverse transcriptase activity.

9. A method of amplifying nucleic acid having complementary nucleotide sequences linked alternately in a single-stranded chain by repeatedly conducting the following steps:

A) the step of providing a template which is provided at the 3'- and 5'-terminals thereof with a region consisting of a nucleotide sequence complementary to each terminal region in the same chain and which upon annealing of these mutually complementary nucleotide sequences, forms a loop capable of base pairing therebetween,

B) the step of performing the synthesis of complementary chain wherein the 3'-terminal of said template annealed to the same chain serves as the origin of synthesis,

C) the step of annealing, to the loop portion, of an oligonucleotide provided at the 3'-terminal thereof with a complementary nucleotide sequence to a loop which among said loops, is located at the 3'-terminal site, followed by synthesis,

with the oligonucleotide as the origin of synthesis, of a complementary chain by a polymerase catalyzing the strand displacement reaction of synthesizing a complementary chain to displace the complementary chain synthesized in step B) to make the 3'-terminal thereof ready for base pairing, and

D) the step wherein the chain with the 3'-terminal made ready for base pairing in step C) serves as a new template.

10. The method according to claim 9, wherein the oligonucleotide in step C) is provided at the 5'-terminal thereof with a nucleotide sequence complementary to the 3'-terminal serving as the origin of synthesis in step B).

11. The method according to claim 10, further comprising the step where a complementary chain synthesized with the oligonucleotide in step C) as the origin of synthesis is used as a template in step A).

12. The method according to claim 9, wherein the template in step A) is synthesized by the method described in claim 5.

13. The method according to claim 1 or 9, wherein the strand displacement reaction of synthesizing complementary chain is carried out in the presence of a melting temperature regulator.

14. The method according to claim 13, wherein the melting temperature regulator is betaine.

15. The method according to claim 14, wherein 0.2 to 3.0 M betaine is allowed to be present in the reaction solution.

16. A method of detecting a target nucleotide sequence in a

sample, which comprises performing an amplification method described in any one of claims 9 to 15 and observing whether an amplification reaction product is generated or not.

17. The method according to claim 16, wherein a probe containing a nucleotide sequence complementary to the loop is added to the amplification reaction product and hybridization therebetween is observed.

18. The method according to claim 17, wherein the probe is labeled on particles and aggregation reaction occurring upon hybridization is observed.

19. The method according to claim 16, wherein an amplification method described in any one of claims 9 to 15 is conducted in the presence of a detector for nucleic acid, and whether an amplification reaction product is generated or not is observed on the basis of a change in the signal of the detector.

20. A method of detecting a mutation in a target nucleotide sequence by the detection method described in claim 16, wherein a mutation in a nucleotide sequence as the subject of amplification prevents synthesis of any one of complementary chains constituting the amplification method.

21. A kit for synthesis of nucleic acid having complementary chains alternately linked in a single-stranded chain, comprising the following elements:

i) the oligonucleotide described in claim 3 wherein the region F2c in nucleic acid as a template is X2c, and F1c located at



the 5'-side of F2c is X1c;

ii) an oligonucleotide containing a nucleotide sequence complementary to an arbitrary region in a complementary chain synthesized with the oligonucleotide in (i) as a primer;

iii) an oligonucleotide having a nucleotide sequence complementary to a region F3c located at the 3'-side of the region F2c in the nucleic acid serving as a template;

iv) a DNA polymerase catalyzing the strand displacement-type reaction of synthesizing complementary chain; and

v) a nucleotide serving as a substrate for the element iv).

22. The kit according to claim 21, wherein the oligonucleotide in ii) is the oligonucleotide described in claim 3 wherein an arbitrary region R2c in a complementary chain synthesized with the oligonucleotide in i) as the origin of synthesis is X2c, and R1c located at the 5' of R2c is X1c.

23. The kit according to claim 22, further comprising:

vi) an oligonucleotide having a nucleotide sequence complementary to a region R3c located at the 3'-side of the arbitrary R2c in a complementary chain synthesized with the oligonucleotide in i) as the origin of synthesis.

24. A kit for detection of a target nucleotide sequence, comprising a detector for detection of a product of nucleic acid synthetic reaction additionally in a kit described in any one of claims 21 to 23.

# ABSTRACT

The present invention relates to an oligonucleotide having a novel structure and a method of synthesizing nucleic acid by using the same as a primer. This oligonucleotide is provided at the 5'-side of the primer with a nucleotide sequence substantially the same as a region synthesized with this primer as the origin of synthesis. The present invention realizes synthesis of nucleic acid based on an isothermal reaction with a simple constitution of reagents. Further, the present invention provides a method of synthesizing highly specific nucleic acid on the basis of this method of synthesizing nucleic acid.

## SEQUENCE LISTING

<110> Eiken Kagaku Kabushiki Kaisha

<120> Method of Synthesizing Nucleic Acid.

<130> E2-001PCT

<140>

<141>

<150> JP-1998-317476

<151> 1998-11-09

<160> 29

<170> PatentIn Ver. 2.0

<210> 1

<211> 52

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 1

cgactctaga ggatccccgg gtactttttg ttgtgtggaa ttgtgagcgg at 52

<210> 2

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 2

acaacgtcgt gactgggaaa accctttttg tgcgggcctc ttcgctatta c 51

<210> 3

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 3

actttatgct tccggctcgt a

21

<210> 4

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 4

gttggaagg gcgatcg

17

<210> 5

<211> 600

<212> DNA

<213> Bacteriophage M13mp18

<400> 5

gcgccaata cgcaaaccgc ctctccccgc gcgttgccg attcattaat gcagctggca 60  
cgacaggttt cccgactgga aagcgggcag tgagcgcaac gcaattaatg tgagttagct 120  
caticattag gcaccccagg ctttacactt tatgcttccg gctcgtagt tgtgtggaat 180  
tgigagcgga taacaatttc acacaggaaa cagctatgac catgattacg aattcgagct 240  
cggatcccgg ggatcctcta gagtcgacct gcaggcatgc aagcttggca ctggccgtcg 300  
ttttacaacg tcgtgactgg gaaaaccctg gcgttaccca acttaatcgc ctgcagcac 360

atcccccttt cgccagctgg cgtaatagcg aagaggcccg caccgatcg ccttcccaac 420  
 agttgcgcag cctgaatggc gaatggcgct ttgcctgggt tccggcacca gaagcgggtgc 480  
 cggaaagctg gctggagtc gatcttcctg aggccgatac ggtcgtcgtc cctcaaaact 540  
 ggcagatgca cggttacgat gcgcccact acaccaacgt aacctatccc attacggtca 600

<210> 6

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
 synthesized primer sequence

<400> 6

ctcttccaaa agtaaggcag gaaatgtgaa accagatcgt aatttggaag acccagcatc 60  
 cag 63

<210> 7

<211> 43

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
 synthesized primer sequence

<400> 7

gtggattcgc actcctcccg ctgacggga cctgcctcgt cgt

43

<210> 8

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 8

gccacctggg tgggaa

16

<210> 9

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 9

ggcgagggag ttctttctct ag

22

<210> 10

<211> 430

<212> DNA

<213> Hepatitis B virus

<400> 10

ctccttgaca ccgctctctgc tctgtatcgg gaggccttag agtctccgga acattgttca 60  
cctcaccata cagcactcag gcaagctatt ctgtgttggg gtgagttaat gaatctggcc 120  
acctgggtgg gaagtaattt ggaagacceca gcatccaggg aattagtagt cagctatgtc 180  
aatgttaata tgggcctaaa aatcagacaa ctattgtggt ttcacatttc ctgccttact 240  
tttgggaagag aaactgtttt ggagtatttg gtaiccttttg gagtgtggat tcgcactcct 300  
cccgcttaca gaccaccaaa tgcacctatc ttatcaaac ttcgggaaac tactgttggt 360  
agacgacgag gcaggteccc tagaagaaga actccctcgc ctgcgagacg aaggtctcaa 420  
tcgccgcgtc 430

<210> 11

<211> 293

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized sequence



<400> 11

```
acaacgtcgt gactgggaaa accctttttg tgcgggcctc ttcgctatta cgccagctgg 60
cgaaaggggg atgtgctgca aggcgattaa gtigggtaac gccagggttt tcccagtcac 120
gacgttgtaa aacgacggcc agtgccaagc ttgcatgcct gcaggtcgac tctagaggat 180
ccccgggtac cgagctcgaa ttcgtaatca tggcatagc tgtttcctgt gtgaaattgt 240
tatccgctca caattccaca caacaaaaag tacccgggga tcctctagag tgc          293
```

<210> 12

<211> 293

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized sequence

<400> 12

```
cgactctaga ggatccccgg gtactttttg ttgtgtggaa ttgtgagcgg ataacaattt 60
cacacaggaa acagctatga ccatgattac gaattcgagc tcggtaccgg gggatcctct 120
agagtcgacc tgcaggcatg caagcttggc actggccgtc gtittacaac gtcgtgactg 180
ggaaaaccct ggcgttacct aacttaatcg ccttcagea catccccctt tcgccagctg 240
gcgtaatagc gaagaggccc gcacaaaaag ggttttccca gtcacgacgt tgt          293
```

<210> 13

<211> 459

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized sequence

<400> 13

acaacgtcgt gactgggaaa accctttttg tgcgggcctc ttcgctatta cgccagctgg 60  
cgaaaggggg atgtgctgca aggcgattaa gttaggtaac gccagggttt tcccagtcac 120  
gacgttgtaa aacgacggcc agtgccaagc ttgcatgcct gcaggtcgac tctagaggat 180  
ccccgggtac cgagctcgaa ttcgtaatca tggatcatagc tgtttcctgt gtgaaattgt 240  
tatccgctca caattccaca caacaaaaag taccgggga tcctctagag tcgacctgca 300  
ggcatgcaag cttaggcactg gccgtcgitt tacaacgtcg tgactgggaa aacctggcg 360  
ttacceact taatcgccit gcagcacatc cccctttcgc cagctggcgt aatagcgaag 420  
aggccgcac aaaaagggtt tcccagtcac cgacgttgt 459

<210> 14

<211> 458

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized sequence

<400> 14

cgactctaga ggatccccgg giactttttg ttgtgtggaa ttgtgagcgg ataacaattt 60  
cacacaggaa acagctatga ccatgattac gaattcgagc tcggtaccgg gggatcctct 120  
agagtgcacc tgcaggcatg caagcttggc aciggccgtc gttttacaac gtcgtgactg 180  
ggaaaaccct ggcgttaccc aacttaatcg ccttgcagca catccccctt tcgccagctg 240  
gcgtaatagc gaagaggccc gcacaaaaag ggttttccca gtcacgacgt tgtaaaacga 300  
cggccagtgc caagcttgca tgcctgcagg tcgactctag aggatccccg ggtaccgagc 360  
tcgaattcgt aatcatggtc atagctgttt cctgtgtgaa attgttatcc gtcacaatt 420  
ccacacaaca aaaagtlacc ggggatccct tagagtcg 458

<210> 15

<211> 790

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized sequence

<400> 15

acaacgtcgt gactgggaaa accctttttg tgcgggcctc ttcgctatta cgccagctgg 60  
cgaaaggggg atgtgctgca aggcgattaa gttgggtaac gccagggttt tcccagtcac 120  
gacgttgtaa aacgacggcc agtgccaagc ttgcatgcct gcaggctgac tctagaggat 180  
ccccgggtac cgagctcgaa ttcgtaatca tggatcatagc tgtttccctgt gtgaaattgt 240  
tatccgctca caattccaca caacaaaaag taccggggga tcccttagag tcgacctgca 300  
ggcatgcaag cttggcactg gccgtcgitt tacaacgtcg tgactgggaa aaccttggcg 360  
ttacceact taatgcctt gcagcacatc cccctttcgc cagctggcgt aatagcgaag 420

aggcccgcac aaaaagggtt ttcccagtcg cgacgttgta aaacgacggc cagtgccaaag 480  
 ctigcatgcc tgcaggtcga ctctagagga tccccgggta ctttttggtg tgtggaattg 540  
 tgagcggata acaatttcac acaggaaaca gctatgacca tgattacgaa ttcgagctcg 600  
 gtacccgggg atcctctaga gtcgacctgc aggcatgcaa gcttggcact ggccgtcggt 660  
 ttacaacgic gtgactggga aaacctggc gttaccaac ttaatcgctt tgcagcacat 720  
 ccccttttcg ccagctggcg taatagcgaa gaggcccgca caaaaagggt ttcccagtc 780  
 acgacgttgt 790

<210> 16

<211> 789

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
 synthesized sequence

<400> 16

cgactctaga ggatccccgg gtactttttg ttgigtggaa ttgtgagcgg ataacaattt 60  
 cacacaggaa acagctatga ccatgattac gaattcgagc tcggtaccgg gggatcctct 120  
 agagtcgacc tgcaggcatg caagcttggc actggccgic gttttacaac gtcgtgactg 180  
 ggaaaacctt ggcgttaccc aacttaatcg ccttgcagca catccccctt tcgccagctg 240  
 gcgtaatagc gaagaggccc gcacaaaaag ggttttccca gtcacgacgt tgtaaaacga 300  
 cggccagtgc caagcttgca tgcctgcagg tcgactctag aggatccccg ggtaccgagc 360  
 tcgaattcgt aatcatggtc atagctgttt ccigtgtgaa attgttatcc gtcacaaatt 420  
 ccacacaaca aaaagtaccc ggggatccct tagagtcgac ctgcaggcat gcaagcttgg 480

cactggccgt cgttttaca cgtcgtgact gggaaaaccc tttttgtgcg ggcctcttcg 540  
 ctattacgcc agctggcgaa agggggaigt gctgcaagge gattaagttg ggtaacgcca 600  
 gggttttccc agtcacgacg ttgtaaaaacg acggccagtg ccaagcttgc atgcctgcag 660  
 gtcgactcta gaggatcccc gggtaccgag ctcgaaatcg taatcatggt catagctgtt 720  
 tcctgtgtga aattgttata cgctcacaat tccacacaac aaaaagtacc cggggatcct 780  
 ctagagtcg 789

<210> 17

<211> 310

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
 synthesized sequence

<400> 17

gtggattcgc actcctcccg ctgacggga cctgcctcgt cgtctaaca cagtagtttc 60  
 cggaagtgtt gataagatag gggcatttgg tggtcigtta gcgggaggag tgcgaatcca 120  
 cactccaaaa gataccaaat actccaaaac agtttcctt ccaaaagtaa ggcaggaaat 180  
 gtgaaaccac aatagttgtc tgatTTTTAG gccatattt acattgacat agctgactac 240  
 taattccctg gatgctgggt ctccaaatt acgatcggg ttacatttc ctgccttact 300  
 tttggaagag 310

<210> 18

<211> 465

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized sequence

<400> 18

```
gtggattcgc actcctcccg ctgacgaggga cctgcctcgt cgtctaacaa cagtagtttc 60
cggaagtgtt gataagatag gggcatttgg tggctctgtaa gcgggaggag tgcgaatcca 120
cactccaaaa gataccaaat actccaaaac agtttctctt caaaagtaa ggcaggaaat 180
gtgaaaccac aatagttgic tgatttttag gcccatatta acattgacat agctgactac 240
taattccctg gatgctgggt ctccaaatt acgatctggt ttcacatttc ctgccttact 300
tttgggaagag aaactgtttt ggagtatttg gtatcttttg gagtgtggat tcgcactcct 360
cccgcttaca gaccacaaa tgcacctatc ttatcaacac ttccggaaac tactgttggt 420
agacgacgag gcaggtcccg atcagcggga ggagtgcgaa tccac 465
```

<210> 19

<211> 20

<212> DNA

<213> M13mp18

<400> 19

ccggggatcc tctagagtcg

20

<210> 20

<213> M13mp18 mutant

ccgggggatcc tctagagtca

20

### ⟨213⟩ Artificial Sequence

<223> Description of Artificial Sequence:Artificially synthesized primer sequence

cgactctaga ggatccccgg tttttgttgt gtggaattgt gagcggat

48

### ⟨213⟩ Artificial Sequence

⟨223⟩ Description of Artificial Sequence:Artificially

synthesized primer sequence

<400> 22

tgactctaga ggatccccgg tttttgttgt gtggaattgt gagcggat

48

<210> 23

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 23

cgtcgtgact gggaaaaccc tttttgtgcg ggcctcttcg ctattac

47

<210> 24

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence



<400> 24

actttaatgct tccggctcgt a

21

<210> 25

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 25

gttggaagg gcgatcg

17

<210> 26

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 26

tggtccatgat gcagtgggca gctttagtct gcggcggtgt tctg

44

<210> 27

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 27

tgctgggtcg gcacagcctg aagctgacct gaaataacctg gcctg

45

<210> 28

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 28

tgcttggtggc ctctcgtg

18

<210> 29

<211> 17

<212> DNA

<213> Artificial Sequence

<220>


<223> Description of Artificial Sequence:Artificially  
synthesized primer sequence

<400> 29

gggtgtggga agctgtg


1 / 18

Fig. 1




2 / 18

Fig. 2




3/18

Fig. 3



4 / 18

Fig. 4



5 / 18

Fig. 5






Fig. 6





Fig. 7

6001 GGGCCCAATA CGCAAACCGC CTCTCCCGC GGGTTGGCCG ATTCATTAAAT GCAGCTGGCA  
6061 CGACAGGTTT CCGACTGGA AAGCGGGCAG TGAGCGCAAC GCAATTAATG TGAGTTAGCT  
6121 CACTCATTAG GCACCCAGG CTTTACACTT TATGCTTCGG GCTCGTATGT TGTGTGGAAT  
6181 TGTGAGGGGA TAACAATTTC ACACAGGAAA CAGCTATGAC CATGATTACG AATTCGAGCT  
6241 CGGTACCCGG GGATCCTCTA GAGTCGACCT GCAGGCATGC AAGCTTGGCA CTGGCOGTGG  
6301 TTTTACAACG TGTGACTGG GAAAACCTG GCGTTACCA ACTTAATGCG CTTCGAGCAC  
6361 ATCCCCCTTT CGCCAGCTGG CGTAATAGCG AAGAGGCGCG CACCGATGCG CCTTCOCAAC  
6421 AGTTGCGCAG CCTGAATGGC GAATGGCGCT TTGCTGGTTC TCGGCACCA GAAGCGGTGC  
6481 CGGAAAGCTG GCTGGAGTGC GATCTTCCTG AGGCGGATAC GGTCGTCGTC CCCTCAAAT  
6541 GGCAGATGCA CGGTTACGAT GCGCCCATCT ACAOCAACGT AACCTATGCC ATTACGGTCA

Diagram illustrating the sequence alignment and primer binding sites for the M13F3, M13F2, M13F1c, M13R1c, M13R2, and M13R3 primers. The sequence is shown in 60-base segments, with the primers indicated by arrows above the corresponding sequence lines.


8 / 18

Fig. 8



9/18


Fig. 9



10/18

Fig. 10

0      0.5      1      2M  
-21 N -21 N -21 N -21 N



11 / 18


Fig. 11

1 CTCCTTGACA CCGCCTCTGC TCTGTATCGG GAGGCTTAG AGTCTCCGGA ACATTGTTCA  
 61 OCTCACCATA CAGCACTCAG GCAAGCTATT CTGTGTTGGG GTGAGTTAAT GAATCTGGOC  
 HBF3 HB65F2  
 121 ACCTGGGTGG GAAGTAATTT GGAAGACCCA GCATCCAGGG AATTAGTAGT CAGCTATGTC  
 HB65F1c  
 181 AATGTTAATA TGGGCTAAA AATCAGACAA CTATTGTGGT TTCACATTC CTGCCCTACT  
 HB65R1c  
 241 TTTGGAAGAG AAACGTGTTT GGAGTATTG GATCTTTTG GAGTGTGGAT TCGCACTCCT  
 301 CCCGCTTACA GACCACCAA TGCCCTATC TTATCAACAC TTCCGGAAAC TACTGTTGTT  
 HB65R2 HBR3  
 361 AGACGACGAG GCAGGTCCCC TAGAAGAAGA ACTCCCTCGC CTCGCAGACG AAGGTCTCAA  
 421 TCGCCGCGTC

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000


12/18

Fig. 12



13 / 18


Fig. 13





14/18

Fig. 14



15 / 18

Fig. 15


6001 GCGCCCAATA CGCAAACCGC CTCTCCCCGC GCGTTGGCCG ATTCATTAAT GCAGCTGGCA  
6061 CGACAGGTTT CCCGACTGGA AAGCGGGCAG TGAGCGCAAC GCAATTAATG TGAGTTAGCT  
6121 CACTCATTAG GCACCCAGG CTTTACACTT TATGCTTCCG GCTCGTATGT TGTGTGGAAT  
6181 TGTGAGCGGA TAACAATTTC ACACAGGAAA CAGCTATGAC CATGATTACG AATTCGAGCT  
6241 CGGTACCCGG GGATCCTCTA GAGTCGACCT GCAGGCATGC AAGCTTGGCA CTGGCCGTCG  
6301 TTTTACAACG TCGTGACTGG GAAAACCCTG GCGTTACCCA ACTTAATCGC CTTGCAGCAC  
6361 ATCCCCCTTT CGCCAGCTGG CGTAATAGCG AAGAGGCCCG CACCGATCGC CCTTCCCAAC  
6421 AGTTGCGCAG CCTGAATGGC GAATGGCGCT TTGCCTGGTT TCCGGCACCA GAAGCGGTGC  
6481 CGGAAAGCTG GCTGGAGTGC GATCTTCCTG AGGCCGATAC GGTCGTCGTC CCCTCAAAC  
6541 GGCAGATGCA CGGTTACGAT GCGCCCATCT ACACCAACGT AACCTATCCC ATTACGGTCA

Diagram illustrating primer binding sites and PCR products for a DNA sequence. The sequence is shown in 10 lines (6001 to 6541). Primers are indicated by arrows above the sequence:

- M13F3 (forward) and M13F2 d4 (reverse) bind to the sequence between 6061 and 6121.
- M13F1c d4 (reverse) binds to the sequence between 6181 and 6241.
- M13R1c d4 (reverse) binds to the sequence between 6241 and 6301.
- M13R2 d4 (reverse) and M13R3 (reverse) bind to the sequence between 6361 and 6421.

A specific nucleotide 'A' is highlighted in the sequence at position 6241 (GAGTCGACCT).

Fig. 16



17/18

Fig. 17

1 ATTCCGCCGG AGAGCTGTGT CACCATGTGG GTCCCGGTTG TCTTCCTCAC CCTGTCCGTG

61 ACGTGGATTG GTGCTGCACC CCTCATCCTG TCTCGGATTG TGGGAGGCTG GGAGTGCGAG

121 AAGCATTTCC AACCCTGGCA GGTGCTTGTG GCCTCTCGTG GCAGGGCAGT CTGCGGCGGT

181 GTTCTGGTGC ACCCCCAGTG GGTCTCACA GCTGCCCACT GCATCAGGAA CAAAAGCGTG

241 ATCTTGCTGG GTCGGCACAG CCTGTTTCAT CCTGAAGACA CAGGCCAGGT ATTTACAGGTC

301 AGCCACAGCT TCCCACACCC GCTCTACGAT ATGAGCCTCC TGAAGAATCG ATTCTCAGG

361 CCAGGTGATG ACTCCAGCCA CGACCTCATG CTGCTCCGCC TGTCAGAGCC TGCCGAGCTC

421 ACGGATGCTG TGAAGGTCAT GGACCTGCCC ACCCAGGAGC CAGCACTGGG GACCACCTGC


481 TACGCCTCAG GCTGGGGCAG CATTGAACCA GAGGAGT

PSAF3 PSAR2 PSAR1c PSAR1c PSAR2 PSAR3

Sau3AI


18/18

Fig. 18



COMBINED DECLARATION FOR PATENT  
APPLICATION AND POWER OF ATTORNEY  
(Includes Reference to PCT International Applications)

ATTORNEY'S DOCKET NUMBER  
201487/1020 (E2-001PCT-US)



As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**METHOD OF SYNTHESIZING NUCLEIC ACID**

the specification of which (check only one item below):

- ☐ is attached hereto.
- ☐ was filed as U.S. Patent Application Serial No. \_\_\_\_\_ on \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable).
- ☒ was filed as PCT International Application Number PCT/JP99/06213 on November 8, 1999 and assigned U.S. Serial Number **09/530,061**.

I hereby state that I have reviewed and understand the contents of the above-identified specifications, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

**PRIOR APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:**

COUNTRY (IF PCT, indicate "PCT")	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119
JP	10/317476	09 November 1998	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
PCT	PCT/JP99/06213	08 November 1999	<input checked="" type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO
			<input type="checkbox"/> YES <input type="checkbox"/> NO

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (Continued) (Includes Reference to PCT International Applications)				ATTORNEY'S DOCKET NUMBER <b>201487/1020 (E2-001PCT-US)</b>	
I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56(a) which occurred between the filing date of the prior application(s) and the national or PCT International filing date of this application:					
PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120:					
U.S. APPLICATIONS			STATUS (Check One)		
U.S. APPLICATION NUMBER	U.S. FILING DATE	PATENTED	PENDING	ABANDONED	
PCT APPLICATIONS DESIGNATING THE U.S.					
PCT APPLICATION NO.	PCT FILING DATE	U.S. SERIAL NUMBERS ASSIGNED (if any)			
PCT/JP99/06213	08 Nov 1999			X	
POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. <b>Michael L. Goldman, Registration No. 30,727; Gunnar G. Leinberg, Registration No. 35,584; Dennis M. Connolly, Registration No. 40,964; Edwin V. Merkel, Registration No. 40,087; Georgia Caton, Registration No. 44,597; Grant E. Pollack, Registration No. 34,097</b>					
Send Correspondence to:			<u>Dennis M. Connolly, Esq.</u> <u>NIXON PEABODY LLP</u> <u>Clinton Square, P.O. Box 1051</u> <u>Rochester, New York 14603</u>		Direct telephone calls to: <b>Dennis M. Connolly</b> <b>(716) 263-1741</b>
1-020	FULL NAME OF INVENTOR	FAMILY NAME <u>Notomi</u>	FIRST GIVEN NAME <u>Tsugunori</u>	SECOND GIVEN NAME	
	RESIDENCE & CITIZENSHIP	CITY <u>Tochigi</u>	STATE/FOREIGN COUNTRY <u>Japan</u> JPX	COUNTRY OF CITIZENSHIP <u>Japan</u>	
	POST OFFICE ADDRESS	P.O. ADDRESS <u>1381-3, Shimoishigami, Otawara-shi</u>	CITY <u>Tochigi</u>	STATE & ZIP CODE/COUNTRY <u>324-0036 Japan</u>	
2-020	FULL NAME OF INVENTOR	FAMILY NAME <u>Hase</u>	FIRST GIVEN NAME <u>Tetsu</u>	SECOND GIVEN NAME	
	RESIDENCE & CITIZENSHIP	CITY <u>Tochigi</u>	STATE/FOREIGN COUNTRY <u>Japan</u> JPX	COUNTRY OF CITIZENSHIP <u>Japan</u>	
	POST OFFICE ADDRESS	P.O. ADDRESS <u>1381-3, Shimoishigami, Otawara-shi</u>	CITY <u>Tochigi</u>	STATE & ZIP CODE/COUNTRY <u>324-0036 Japan</u>	
2-033	FULL NAME OF INVENTOR	FAMILY NAME	FIRST GIVEN NAME	SECOND GIVEN NAME	
	RESIDENCE & CITIZENSHIP	CITY	STATE/FOREIGN COUNTRY	COUNTRY OF CITIZENSHIP	
	POST OFFICE ADDRESS	P.O. ADDRESS	CITY	STATE & ZIP CODE/COUNTRY	
I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.					
SIGNATURE OF INVENTOR 201 <u>T. Notomi</u>		SIGNATURE OF INVENTOR 202 <u>Tetsu Hase</u>		SIGNATURE OF INVENTOR 203	
DATE <u>14. June 2000</u>		DATE <u>18 June 2000</u>		DATE	